

FORMALITY THEOREMS FOR HOCHSCHILD CHAINS IN THE LIE ALGEBROID SETTING

DAMIEN CALAQUE, VASILIIY DOLGUSHEV AND GILLES HALBOUT

ABSTRACT. In this paper we prove Lie algebroid versions of Tsygan's formality conjecture for Hochschild chains both in the smooth and holomorphic settings. Our result in the holomorphic setting implies a version of Tsygan's formality conjecture for Hochschild chains of the structure sheaf of any complex manifold. The proofs are based on the use of Kontsevich's quasi-isomorphism for Hochschild cochains of $\mathbb{R}[[y^1, \dots, y^d]]$, Shoikhet's quasi-isomorphism for Hochschild chains of $\mathbb{R}[[y^1, \dots, y^d]]$, and Fedosov's resolutions of the natural analogues of Hochschild (co)chain complexes associated with a Lie algebroid. In the smooth setting we discuss an application of our result to the description of quantum traces for a Poisson Lie algebroid.

CONTENTS

Introduction	2
Notations	3
Acknowledgements	3
1. Algebraic structures associated with a Lie algebroid	3
1.1. Lie algebroids and associated sheaves	3
1.2. Lie algebroids connections	6
1.3. Algebraic structures on E -polydifferential operators and E -polyjets	7
2. The formality theorem for E -chains	11
2.1. Hochschild-Kostant-Rosenberg	11
2.2. The formality of the DGLA module of E -chains	12
2.3. Formality theorems for the Hochschild complexes of $\mathbb{R}[[y^1, \dots, y^d]]$.	13
3. The Fedosov resolutions	14
4. Proof of the formality theorem for E -chains and its applications	25
4.1. Proof of the theorem	25
4.2. Applications of the formality theorem	28
5. Formality theorems for holomorphic Lie algebroids	30
5.1. Algebraic structures and the main theorem	30
5.2. Fedosov resolutions	31
5.3. End of the proof of theorem 5.2	37
6. Concluding remarks	39
References	39

INTRODUCTION

Lie algebroids and Lie groupoids provide a natural framework for developing analysis on differentiable foliations and manifolds with corners [24], [25], [27], [38]. This motivates our interest to the natural analogues of Hochschild and cyclic (co)homological complexes in the setting of Lie algebroids and to the corresponding analogues of the Kontsevich-Tsygan formality conjectures. Thus the formality theorem for the differential graded Lie algebra (DGLA) of Hochschild cochains in the Lie algebroid setting [1] allows us to quantize an arbitrary Poisson Lie algebroid¹. The formality of the DGLA module of Hochschild chains in the Lie algebroid setting would give a description of the quantum traces for Poisson Lie algebroids, and the formality of the cyclic complex in the setting of Lie algebroids would imply the algebraic index theorem [26], [32] for the deformations associated with an arbitrary Poisson Lie algebroid.

An appropriate analogue of the Hochschild cochain (resp. chain) complex associated with a Lie algebroid E is the complex of E -polydifferential operators (resp. Hochschild E -chains) (see definitions 1.9 and 1.14 in the next section). It turns out that the complex of E -polydifferential operators is naturally a DGLA and the complex of E -chains is naturally a DG module over this DGLA. Due to the recent result [1] of the first author for any Lie algebroid E over a smooth manifold the DGLA of E -polydifferential operators is formal.

In this paper we use Kontsevich's [20] and Shoikhet's [29] formality theorems for \mathbb{R}_{formal}^d and the 'Fedosov-like' [12] globalization technique [3, 9, 10, 26] to prove that for any Lie algebroid E over a smooth manifold (resp. holomorphic Lie algebroid over a complex manifold) the DGLA module of E -chains (resp. the sheaf of DGLA modules of E -chains) is formal. In the smooth setting this result allows us to describe quantum traces for an arbitrary Poisson Lie algebroid. In the holomorphic setting this result implies a version of Tsygan's formality conjecture for Hochschild chains of the structure sheaf of any complex manifold.

Eliminating the sheaf of Hochschild E -chains in the holomorphic setting we get that for any holomorphic Lie algebroid E the sheaf of E -polydifferential operators is formal as a sheaf of DGLAs. In particular, this result implies Kontsevich's formality theorem for complex manifolds, the proof of which was formulated only for algebraic varieties [39].

The paper is organized as follows. In the first section we recall some basic facts about Lie algebroids and define algebraic structures on the complexes of E -polydifferential operators and E -polyjets of an algebroid E . We recall Kontsevich's [20] and Shoikhet's [29] formality theorems for \mathbb{R}_{formal}^d and formulate our first result, the formality of the module of E -chains (see theorem 2.2 on page 12). The second section is devoted to the construction of the Fedosov resolutions of the sheaves of E -polydifferential operators, E -chains, E -polyvector fields and E -forms. It is the most technical part of the paper. Using these resolutions in section 3, we prove theorem 2.2. In the same section we apply this theorem to the description of quantum traces of Poisson Lie algebroids. In section 4 we prove Tsygan's formality conjecture for Lie algebroid chains in the holomorphic setting (see theorem 5.2 on page 31), which, in particular, gives us the formality theorem for Hochschild chains

¹According to the terminology of P. Xu [37] we have to call this object a triangular Lie bialgebroid. However, since we do not mention the bialgebroid structure, we refer to this object as a Poisson Lie algebroid.

of the structure sheaf of an arbitrary complex manifold (see theorem 5.4) In the concluding section we mention an equivariant version of theorem 2.2 and raise some other questions.

Notations. We assume Einstein’s convention for the summation over repeated indices and omit the symbol \wedge referring to a local basis of exterior forms. The arrow \succrightarrow denotes an L_∞ -morphism of L_∞ -algebras, the arrow $\succ\succrightarrow$ denotes a morphism of L_∞ -modules, and the notation

$$\begin{array}{c} \mathcal{L} \\ \downarrow_{mod} \\ \mathcal{M} \end{array}$$

means that \mathcal{M} is an L_∞ -module over the L_∞ -algebra \mathcal{L} . The abbreviation “DGLA” stands for “differential graded Lie algebra” and the abbreviation “DGA” stands for “differential graded associative algebra”. Throughout the paper (except section 5) we work over the field \mathbb{R} of real numbers: unless otherwise specified, M denotes a smooth real manifold, \mathcal{O}_M denotes the sheaf of real valued C^∞ -functions on M and vector bundles are real vector bundles. Finally, we denote by the same symbol a vector bundle and its sheaf of sections.

Acknowledgements. We would like to thank G. Felder and A. Cattaneo for their interest to this work. The second author is partially supported by the NSF grant DMS-9988796, the Grant for Support of Scientific Schools NSh-1999.2003.2 and the grant CRDF RM1-2545-MO-03.

1. ALGEBRAIC STRUCTURES ASSOCIATED WITH A LIE ALGEBROID

1.1. **Lie algebroids and associated sheaves.** Let us recall the following

Definition 1.1. *A Lie algebroid over a smooth manifold M is a smooth vector bundle E of finite rank whose sheaf of sections is a sheaf of Lie algebras equipped with a \mathcal{O}_M -linear morphism of sheaves of Lie algebras*

$$\rho : E \rightarrow TM .$$

The \mathcal{O}_M -module structure and the Lie algebra structure on the sheaf E are compatible in the following sense: for any open subset $U \subset M$, any function $f \in \mathcal{O}_M(U)$ and any sections $u, v \in \Gamma(U, E)$

$$(1.1) \quad [u, fv] = f[u, v] + \rho(u)(f)v .$$

The map ρ is called the anchor.

Examples. 1. The tangent bundle TM on M is the simplest example of a Lie algebroid. The bracket is the usual Lie bracket of vector fields and the anchor is the identity map $\text{id} : TM \rightarrow TM$.

2. More generally any involutive distribution (i.e. regular foliation) $E \subset TM$ is a Lie algebroid over M .

3. A Lie algebroid over a point is simply a finite dimensional Lie algebra.

1.1.1. *The sheaf of E -polyvector fields.*

Definition 1.2. *The bundle ${}^E T_{poly}^*$ of E -polyvector fields is the exterior algebra of the bundle E with the shifted grading*

$$(1.2) \quad {}^E T_{poly}^* = \bigoplus_{k \geq -1} {}^E T_{poly}^k, \quad {}^E T_{poly}^k := \wedge^{k+1} E.$$

It turns out that the Lie bracket $[\cdot, \cdot]$ on sections of ${}^E T_{poly}^0 = E$ can be naturally extended to a Lie bracket on sections of the whole vector bundle ${}^E T_{poly}^*$ of E -polyvectors (it was noticed in [2]). Indeed, first, we defined a Lie bracket $[\cdot, \cdot]_{SN}$ on homogeneous sections of low degree as follows:

$$(1.3) \quad [f, g]_{SN} := 0, \quad [u, f]_{SN} := \rho(u)f, \quad \text{and} \quad [u, v]_{SN} := [u, v].$$

$$\forall f, g \in \mathcal{O}_M(U), \quad u, v \in \Gamma(U, E)$$

Next, we extend $[\cdot, \cdot]_{SN}$ to sections of ${}^E T_{poly}^*$ (i.e. E -polyvector fields) by requiring the graded Leibniz rule with respect to the \wedge -product:

$$(1.4) \quad [u, v \wedge w]_{SN} = [u, v]_{SN} \wedge w + (-1)^{k(l+1)} v \wedge [u, w]_{SN},$$

$$\forall u \in \Gamma(U, {}^E T_{poly}^k), \quad v \in \Gamma(U, {}^E T_{poly}^l), \quad w \in \Gamma(U, {}^E T_{poly}^*).$$

In the simplest example $E = TM$ the Lie bracket $[\cdot, \cdot]_{SN}$ coincides with the well known Schouten-Nijenhuis bracket of ordinary polyvector fields.

1.1.2. *The sheaf of E -differential forms.*

The exterior algebra $\wedge^* E^\vee$ of the dual bundle E^\vee to E is a natural candidate for the bundle ${}^E \Omega_M^*$ of E -differential forms. Sections of ${}^E \Omega_M^*$ (E -forms for short) are endowed with the following E -de Rham differential

$$(1.5) \quad {}^E d\omega(\sigma_0, \dots, \sigma_k) := \sum_i (-1)^i \rho(\sigma_i) \omega(\sigma_0, \dots, \hat{\sigma}_i, \dots, \sigma_k)$$

$$+ \sum_{i < j} (-1)^{i+j} \omega([\sigma_i, \sigma_j], \sigma_0, \dots, \hat{\sigma}_i, \dots, \hat{\sigma}_j, \dots, \sigma_k),$$

$$\sigma_i \in \Gamma(U, E).$$

Another operation defined on E -forms is the contraction with E -polyvector fields. For a E -polyvector field $u \in \Gamma(U, {}^E T_{poly}^k)$ we denote by ι_u the contraction with u . Using this contraction, the E -de Rham differential (1.5), and the Cartan-Weil formula

$$(1.6) \quad {}^E L_u := {}^E d \circ \iota_u + (-1)^k \iota_u \circ {}^E d$$

we define the E -Lie derivative of E -forms (over an open subset U) by the E -polyvector field $u \in \Gamma(U, {}^E T_{poly}^k)$.

For our purposes it is more convenient to use the reversed grading in the bundle of E -forms. Thus we denote by

$$(1.7) \quad {}^E A_* = {}^E \Omega_M^{-*}$$

the corresponding bundle with reversed grading and observe that ${}^E A_*$ is equipped with a structure of a graded module over the sheaf of graded Lie algebras ${}^E T_{poly}^*$ via the E -Lie derivative (1.6). Namely,

Lemma 1.3. *For any $u \in \Gamma(U, {}^E T_{poly}^k)$ and $v \in \Gamma(U, {}^E T_{poly}^l)$ one has*

$$(1.8) \quad {}^E L_u \circ {}^E L_v - (-1)^{kl} {}^E L_v \circ {}^E L_u = {}^E L_{[u,v]_{SN}}.$$

Proof. First, it is immediate from the definition (1.6) that for any $u \in \Gamma(U, {}^E T_{poly}^k)$

$$(1.9) \quad {}^E d \circ {}^E L_u = (-1)^k {}^E L_u \circ {}^E d.$$

Second, we claim that for any $v \in \Gamma(U, {}^E T_{poly}^l)$ we have

$$(1.10) \quad {}^E L_u \circ \iota_v - (-1)^{k(l+1)} \iota_v \circ {}^E L_u = (-1)^k \iota_{[u,v]_{SN}}.$$

Using (1.9) and (1.10) it is not hard to show that

$${}^E L_u ({}^E d \iota_v + (-1)^l \iota_v {}^E d) - (-1)^{kl} ({}^E d \iota_v + (-1)^l \iota_v {}^E d) {}^E L_u = ({}^E d \iota_{[u,v]_{SN}} + (-1)^{k+l} \iota_{[u,v]_{SN}} {}^E d).$$

Thus it suffices to prove that equation (1.10) holds.

The proof of (1.10) goes as follows. First, direct computations show that (1.10) holds for any sections u and v of the subsheaf ${}^E T_{poly}^{-1} \oplus {}^E T_{poly}^0$. Second, using the Leibniz rule (1.4) we prove the desired identity by induction on the degrees of E -polyvector fields u and v . In doing this, we need another simple identity

$${}^E L_{u_1 \wedge u_2} = {}^E L_{u_1} \iota_{u_2} - (-1)^{k_1} \iota_{u_1} {}^E L_{u_2}, \quad \forall u_i \in \Gamma(U, {}^E T_{poly}^{k_i}),$$

which follows easily from the fact that $\iota_{u_1 \wedge u_2} = \iota_{u_1} \circ \iota_{u_2}$. \square

1.1.3. The sheaf of E -differential operators.

One can also define the (left) \mathcal{O}_M -module $\mathcal{U}E$ of E -differential operators to be the sheaf of algebras locally generated by functions and E -vector fields. More precisely, $\mathcal{U}E$ is the sheaf associated with the following presheaf

$$(1.11) \quad U \longmapsto \frac{T(\mathcal{O}_M(U) \oplus \Gamma(U, E))}{\left\{ \begin{array}{l} f \otimes g - fg, f \otimes u - fu, \\ u \otimes f - f \otimes u - \rho(u)f, \\ u \otimes v - v \otimes u - [u, v], \end{array} \right\}}$$

$$f, g \in \mathcal{O}_M(U), \quad u, v \in \Gamma(U, E).$$

As a sheaf of \mathcal{O}_M -modules, $\mathcal{U}E$ is endowed with an increasing filtration

$$(1.12) \quad \mathcal{O}_M = \mathcal{U}E^0 \subset \mathcal{U}E^1 \subset \mathcal{U}E^2 \subset \dots \subset \mathcal{U}E,$$

which is defined by assigning the degree 1 to the E -polyvector fields.

In the terminology of [28] E is a sheaf of *Lie-Rinehart algebras* over the structure sheaf \mathcal{O}_M and $\mathcal{U}E$ is its universal enveloping algebra. Besides the fact that $\mathcal{U}E$ is a sheaf of algebras, $\mathcal{U}E$ is also equipped with a coassociative \mathcal{O}_M -linear map $\Delta : \mathcal{U}E \rightarrow \mathcal{U}E \otimes_{\mathcal{O}_M} \mathcal{U}E$ which is defined as follows

$$\Delta(f) = f \otimes 1 = 1 \otimes f,$$

$$(1.13) \quad \Delta(u) = u \otimes 1 + 1 \otimes u, \quad \Delta(PQ) = \Delta(P)\Delta(Q),$$

$$\forall u \in \Gamma(U, E), \quad P, Q \in \Gamma(U, \mathcal{U}E).$$

Remark. For any $u \in \Gamma(U, \mathcal{O}_M \oplus E)$ one can see that a lift of $\Delta(u)$ lies in the normalizer $N(\mathcal{I}_U)$ of the right ideal \mathcal{I}_U generated by $f \otimes 1 - 1 \otimes f$, $f \in \mathcal{O}_M(U)$, in $\Gamma(U, \mathcal{U}E \otimes_{\mathbb{R}} \mathcal{U}E)$. Therefore Δ takes values in a sheaf of algebras (the one associated

to the presheaf of algebras $U \mapsto N(\mathcal{I}_U)/\mathcal{I}_U$; hence $\Delta(PQ) = \Delta(P)\Delta(Q)$ is well-defined.

Moreover the anchor ρ extends to a (left) \mathcal{O}_M -linear morphism of sheaves of (associative) algebras $\rho : \mathcal{U}E \rightarrow \text{End}(\mathcal{O}_M)$. In the terminology of [37] $(\mathcal{U}E, \Delta, \rho)$ is a sheaf of *Hopf algebroids with anchor*.

Notice that, in the simplest example $E = TM$ of the Lie algebroid $\mathcal{U}E$ is the sheaf of differential operators on M . In this case $\Delta(P)$ is the bidifferential operator $(f, g) \mapsto P(fg)$.

The following result shows that $\mathcal{U}E$ is an ind-finite dimensional vector bundle over M .

Proposition 1.4 ([27, 28]). *$\mathcal{U}E \cong S(E)$ as sheaves of (left) \mathcal{O}_M -modules.*

1.2. Lie algebroids connections. By the word *connection* on a vector bundle \mathcal{B} over M we always mean *E-connection*, that is a linear operator

$$(1.14) \quad \nabla : \Gamma(M, \mathcal{B}) \rightarrow {}^E\Omega^1(M, \mathcal{B})$$

satisfying the following equation

$$(1.15) \quad \nabla(fu) = {}^E d(f)u + f\nabla(u)$$

for any $f \in \mathcal{O}_M(M)$ and $u \in \Gamma(M, \mathcal{B})$.

Locally, ∇ is completely determined by its *Christophel's symbols* Γ_{ij}^k . Namely, let (e_1, \dots, e_r) and (ξ^1, \dots, ξ^r) be dual local basis of E and E^\vee respectively, and (b_1, \dots, b_s) be a local base of \mathcal{B} , then

$$(1.16) \quad \nabla(b_j) = \xi^i \Gamma_{ij}^k b_k$$

For any $u \in \Gamma(M, E)$ we denote by ∇_u the associated map $\Gamma(M, \mathcal{B}) \rightarrow \Gamma(M, \mathcal{B})$.

Remark. As with usual connections, one can extend this *covariant derivative* on E -tensor in a unique way such that ∇_u is a derivation with respect to the tensor product of E -tensors, commutes with the contraction of E -tensors, acts as $\rho(u)$ on functions, and is \mathbb{R} -linear.

Definition 1.5. *The curvature R of a connection ∇ with value in \mathcal{B} is the section R of the bundle $E^\vee \otimes E^\vee \otimes \mathcal{B}^\vee \otimes \mathcal{B}$ defined by*

$$(1.17) \quad R(u, v)w = (\nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{[u, v]})w$$

for any $u, v \in \Gamma(M, E)$ and $w \in \Gamma(M, \mathcal{B})$.

Locally, the curvature is given by

$$R(e_i, e_j)b_k = (R_{ij})_k^l b_l$$

with

$$(1.18) \quad (R_{ij})_k^l = \Gamma_{im}^l \Gamma_{jk}^m - \Gamma_{ik}^m \Gamma_{jm}^l + \rho(e_i) \cdot \Gamma_{jk}^l - \rho(e_j) \cdot \Gamma_{ik}^l - c_{ij}^m \Gamma_{mk}^l$$

For a connection ∇ on E itself one has the following

Definition 1.6. *The torsion T of ∇ is a E -tensor of type $(1, 2)$ defined by*

$$(1.19) \quad T(u, v) = \nabla_u v - \nabla_v u - [u, v]$$

for any $u, v \in \Gamma(M, E)$.

One can write the local coefficients of this tensor very easily:

$$(1.20) \quad T_{ij}^k = \Gamma_{ij}^k - \Gamma_{ji}^k - c_{ij}^k$$

Proposition 1.7. *A torsion free connection on E exists.*

Proof. Let $(U_\alpha)_\alpha$ be a cover of M by trivializing opens for E . On each U_α one has a base $(e_i)_i$ of sections and then can define $\nabla_{e_i}^{(\alpha)} e_j = \frac{1}{2}[e_i, e_j]$. Let $(f_\alpha)_\alpha$ be partition of unity for $(U_\alpha)_\alpha$ and define $\nabla = f_\alpha \nabla^{(\alpha)}$. ∇ is a torsion free connection on E . \square

Proposition 1.8 (Bianchi's identities). *Let ∇ be connection on E . For any $u, v, w \in \Gamma(M, E)$ one has*

$$(1.21) \quad \nabla_u R(v, w) + R(T(u, v), w) + c.p.(u, v, w) = 0$$

and

$$(1.22) \quad R(u, v)w - T(T(u, v), w) - \nabla_u T(v, w) + c.p.(u, v, w) = 0$$

Proof. See for example [15]. \square

1.3. Algebraic structures on E -polydifferential operators and E -polyjets.

1.3.1. The sheaf of E -polydifferential operators.

Definition 1.9. *The (ind-finite dimensional) graded bundle ${}^E D_{poly}^*$ of E -polydifferential operators is the tensor algebra of the bundle $\mathcal{U}E$ with a shifted grading:*

$${}^E D_{poly}^* = \bigoplus_{k \geq -1} {}^E D_{poly}^k, \quad {}^E D_{poly}^k = \otimes_{\mathcal{O}_M}^{k+1} \mathcal{U}E.$$

It is easy to see that in the case $E = TM$ the sheaf ${}^E D_{poly}^*$ is the sheaf of polydifferential operators on M .

Using the coproduct (1.13) in $\mathcal{U}E$ we endow the graded sheaf ${}^E D_{poly}^*$ of E -polydifferential operators with a Lie bracket $[\cdot]_G$. To introduce this bracket we first define the following bilinear product of degree 0

$$\bullet : {}^E D_{poly} \otimes {}^E D_{poly} \rightarrow {}^E D_{poly},$$

$$(1.23) \quad \begin{aligned} P \bullet Q &= \sum_{i=0}^{|P|} (-1)^{i|Q|} (1^{\otimes i} \otimes \Delta^{(|Q|)} \otimes 1^{\otimes |P|-i})(P) \cdot (1^{\otimes i} \otimes Q \otimes 1^{\otimes |P|-i}), \\ P \bullet f &= \sum_{i=0}^{|P|} (-1)^i (1^{\otimes i} \otimes \rho \otimes 1^{\otimes |P|-i})(P) (1^{\otimes i} \otimes f \otimes 1^{\otimes |P|-i}), \\ f \bullet g &= 0, \quad f \bullet P = 0, \end{aligned}$$

for any $P, Q \in \Gamma(U, {}^E D_{poly}^{\geq 0})$ and $f, g \in \Gamma(U, {}^E D_{poly}^{-1}) = \mathcal{O}_M(U)$. Here $\Delta^{(n)} = (\Delta \otimes 1^{\otimes n-1}) \circ \dots \circ \Delta$, $\Delta^{(0)}$ is by convention the identity map.

Remark. Let \mathcal{I} be the right ideal in $\mathcal{U}E^{\otimes k}$ generated by $1^{\otimes i-1} \otimes f \otimes 1^{\otimes k-i} - 1^{\otimes i} \otimes f \otimes 1^{\otimes k-i-1}$, where $i = 1, \dots, k-1$ and $f \in \mathcal{O}_M$. The sheaf of algebras $N(\mathcal{I})/\mathcal{I}$ acts on any tensor product $V_1 \otimes_{\mathcal{O}_M} \dots \otimes_{\mathcal{O}_M} V_k$ over \mathcal{O}_M of left $\mathcal{U}E$ -modules V_i 's. Since $\Delta^{(r-1)}$ obviously takes values in $N(\mathcal{I})/\mathcal{I}$ then equation (1.23) is well-defined.

Although the bilinear product is not associative, the graded commutator

$$(1.24) \quad [P, Q]_G = P \bullet Q - (-1)^{|P||Q|} Q \bullet P, \quad P, Q \in \Gamma(U, {}^E D_{poly}^*).$$

defines a graded Lie bracket between the E -polydifferential operators.

It is not hard to see that in the case $E = TM$ the above bracket reduces to the well known Gerstenhaber bracket [16] between polydifferential operators on M .

Notice that an element $1 \otimes 1 \in \Gamma(M, {}^E D_{poly}^1)$ is distinguished by the following remarkable identity $[1 \otimes 1, 1 \otimes 1]_G = 0$. Using this observation we define the following differential

$$(1.25) \quad \partial = [1 \otimes 1, \]_G : {}^E D_{poly}^* \rightarrow {}^E D_{poly}^{*+1}$$

on the sheaf of E -polydifferential operators.

By definition we see that ∂ is compatible with the Lie bracket (1.24). Thus, $({}^E D_{poly}^*, \partial, [\cdot, \cdot]_G)$ is a sheaf of differential graded Lie algebras (DGLA for short).

We would like to mention that the tensor product of sections (over \mathcal{O}_M) turns the sheaf ${}^E D_{poly}[-1]^*$ with the shifted grading into a sheaf of graded associative algebras. Moreover, it is not hard to see that the differential ∂ (1.25) is compatible with this product. Thus ${}^E D_{poly}^*$ can be also viewed as a sheaf of DG associative algebras (DGA).

Remark. Notice that this construction works not only for $\mathcal{U}E$ but for any (sheaf of) Hopf algebroid with anchor. Below, we use the fact that any morphism of Hopf algebroids with anchor induces a morphism between the corresponding DGLAs (resp. DGAs).

1.3.2. The sheaf of E -polyjets.

Definition 1.10. *The bundle EJ_*^{poly} of E -polyjets is the following graded bundle placed in nonnegative degrees*

$$EJ_*^{poly} = \bigoplus_{k \geq 0} EJ_k^{poly}, \quad EJ_k^{poly} := Hom_{\mathcal{O}_M}(\mathcal{U}E^{\otimes_{\mathcal{O}_M} k+1}, \mathcal{O}_M).$$

Since the sheaf ${}^E D_{poly}^*$ of E -polydifferential operators is an ind-finite dimensional graded vector bundle the sheaf EJ_*^{poly} of E -polyjets is a profinite dimensional graded vector bundle. Furthermore, the sheaf EJ_*^{poly} is endowed with a canonical flat connection ∇^G which is called the *Grothendieck connection* and defined by the formula

$$(1.26) \quad \nabla_\sigma^G(j)(P) := \rho(\sigma)(j(P)) - j(\sigma \bullet P),$$

where $\sigma \in \Gamma(U, E)$, $j \in \Gamma(U, EJ_k^{poly})$, $P \in \Gamma(U, {}^E D_{poly}^k)$, and the operation \bullet is defined in (1.23).

For this connection we have the following standard

Proposition 1.11. *Let χ be a map of sheaves*

$$\chi : EJ_k^{poly} \rightarrow \begin{cases} EJ_{k-1}^{poly}, & \text{if } k > 0, \\ \mathcal{O}_M, & \text{if } k = 0 \end{cases}$$

defined by the formula

$$(1.27) \quad \chi(a)(P) = a(1 \otimes P), \quad P \in \Gamma(U, {}^E D_{poly}^{k-1}), \quad a \in \Gamma(U, EJ_k^{poly}).$$

The restriction of the map χ to the ∇^G -flat E -polyjets gives the isomorphism of sheaves

$$(1.28) \quad \chi : \ker \nabla^G \cap EJ_k^{poly} \simeq \begin{cases} EJ_{k-1}^{poly}, & \text{if } k > 0, \\ \mathcal{O}_M, & \text{if } k = 0. \end{cases}$$

Proof. To see that the map (1.28) is surjective one has to notice that for any E -polyjet b of degree $k-1$ (resp. any function b) the equations

$$a(1 \otimes P) = b(P), \quad P \in \Gamma(U, ED_{poly}^{k-1})$$

and

$$(1.29) \quad \begin{aligned} a(u \cdot Q \otimes P) &= \rho(u)a(Q \otimes P) - a(Q \otimes (\Delta^{(k-1)}(u) \cdot P)), \\ Q &\in \Gamma(U, \mathcal{U}E), \quad u \in \Gamma(U, E) \end{aligned}$$

define a ∇^G -flat E -polyjet a of degree k (resp. a ∇^G -flat E -jet a).

On the other hand, if a is a ∇^G -flat E -polyjet of degree k equation (1.29) is automatically satisfied. Thus a is uniquely determined by its image $\chi(a)$. \square

Let t be the cyclic permutation acting on the sheaf EJ_*^{poly} of E -polyjets

$$(1.30) \quad \begin{aligned} t(a)(P_0 \otimes \cdots \otimes P_l) &:= a(P_1 \otimes \cdots \otimes P_l \otimes P_0), \\ a &\in \Gamma(U, EJ_l^{poly}), \quad P_i \in \Gamma(U, \mathcal{U}E). \end{aligned}$$

Using this operation and the bilinear product (1.23) we define the map

$$\begin{aligned} {}^E S : ED_{poly}^k \otimes EJ_l^{poly} &\rightarrow EJ_{l-k}^{poly}, \\ P \otimes a &\mapsto {}^E S_P(a) \end{aligned}$$

such that for $P \in \Gamma(U, ED_{poly}^k)$, $a \in \Gamma(U, EJ_l^{poly})$, and $Q \in \Gamma(U, ED_{poly}^{l-k})$,

$$(1.31) \quad {}^E S_P(a)(Q) = a(Q \bullet P) + \sum_{j=1}^k (-1)^{lj} t^j(a)((\Delta^{(k)} \otimes 1^{\otimes (l-k)})(Q) \cdot (P \otimes 1^{\otimes (l-k)})).$$

Due to the following proposition the map ${}^E S$ defines an action of the sheaf of graded Lie algebras ED_{poly}^* of E -polydifferential operators on the graded sheaf EJ_*^{poly} of E -polyjets. Namely,

Proposition 1.12. *For any pair $P_1, P_2 \in \Gamma(U, ED_{poly}^*)$ of E -polydifferential operators and any E -polyjet $a \in \Gamma(U, EJ_*^{poly})$*

$$(1.32) \quad {}^E S_{P_1} {}^E S_{P_2}(a) - (-1)^{|P_1||P_2|} {}^E S_{P_2} {}^E S_{P_1}(a) = {}^E S_{[P_1, P_2]_G}(a).$$

Moreover, the action (1.31) is compatible with the Grothendieck connection (1.26)

$$(1.33) \quad \nabla_u^G ({}^E S_{P_1}(a)) = {}^E S_{P_1}(\nabla_u^G(a)), \quad u \in \Gamma(U, E).$$

Proof. It is not hard to show that

$$(1.34) \quad {}^E S_{P_1} {}^E S_{P_2}(a) = {}^E S_{P_1 \bullet P_2}(a) + H(P_1, P_2)(a) + (-1)^{|P_1||P_2|} H(P_2, P_1)(a),$$

where²

$$H(P_1, P_2) : EJ_*^{poly} \rightarrow EJ_{* - |P_1| - |P_2|}^{poly}$$

²Formula (1.34) is essentially borrowed from paper [17] of E. Getzler.

is a graded \mathcal{O}_M -linear endomorphism of the sheaf EJ_*^{poly} defined by the following formula

$$\begin{aligned} (H(P_1, P_2)(a))(Q) = & \\ \sum_{i,j} (-1)^{i|P_1|+j|P_2|} a & \left[(1^{\otimes i} \otimes \Delta^{|P_1|} \otimes 1^{\otimes(j-i-|P_1|-1)} \otimes \Delta^{|P_2|} \otimes 1^{\otimes(n-j-|P_2|)}(Q)) \right. \\ & \left. (1^{\otimes i} \otimes P_1 \otimes 1^{\otimes(j-i-|P_1|-1)} \otimes P_2 \otimes 1^{\otimes(n-j-|P_2|)}) \right] + \\ \sum_{k,l} (-1)^{k|P_2|+l(n-|P_2|)} t^l(a) & \left[(\Delta^{|P_1|} \otimes 1^{k+l-|P_1|-1} \otimes \Delta^{|P_2|} \otimes 1^{\otimes n-k-l-|P_2|} (Q)) \right. \\ & \left. P_1 \otimes 1^{\otimes(k+l-|P_1|-1)} \otimes P_2 \otimes 1^{\otimes(n-k-l-|P_2|)} \right], \end{aligned}$$

the sums run over all i, j, k, l satisfying the conditions

$$\begin{aligned} 0 \leq i \leq j - |P_1| - 1, \quad j \leq n - |P_2|, \\ 1 \leq l \leq |P_1|, \quad |P_1| - l + 1 \leq k \leq n - |P_2| - l, \end{aligned}$$

and

$$Q \in \Gamma(U, ED_{poly}^{n-|P_1|-|P_2|}).$$

Equation (1.34) obviously implies identity (1.32).

Equation (1.33) follows immediately from the fact that the coproduct (1.13) is compatible with the multiplication of the E -differential operators and the fact that the Grothendieck connection (1.26) commutes with the cyclic permutation (1.30). \square

Like in the case of E -polydifferential operators the element $1 \otimes 1 \in \Gamma(M, ED_{poly}^1)$ is (satisfying $[1 \otimes 1, 1 \otimes 1]_G = 0$) allows us to define the following differential

$$(1.35) \quad \mathfrak{b} := E_{S_{1 \otimes 1}}^\alpha : EJ_*^{poly} \rightarrow EJ_{*-1}^{poly}$$

on the sheaf of E -polyjets.

From the definition of the differentials (1.25), (1.35) and equation (1.32), we see that \mathfrak{b} is compatible with the action (1.31) in the sense of the following equation

$$\begin{aligned} \mathfrak{b}(E S_P(a)) &= E S_{\partial P}(a) + (-)^{|P|} E S_P(\mathfrak{b}(a)). \\ \forall a \in \Gamma(U, EJ_*^{poly}), \quad P &\in \Gamma(U, ED_{poly}^* \otimes \mathcal{O}_M). \end{aligned}$$

Thus, $(EJ_*^{poly}, \mathfrak{b}, ES)$ is a sheaf of differential graded modules (DG modules for short) over ED_{poly}^* .

1.3.3. Hochschild E -chains.

The complex of sheaves $(EJ_*^{poly}, \mathfrak{b})$ is not a good candidate for the Hochschild chain complex in the Lie algebroid setting. Indeed, if our Lie algebroid E is TM then the complex $(EJ_*^{poly}, \mathfrak{b})$ boils down to the Hochschild chain complex of \mathcal{O}_M without the zeroth term and the action (1.31) does not coincide with the standard action of Hochschild cochains on Hochschild chains (see eq. (3.4) in [10]). To cure these problems simultaneously we introduce a graded sheaf EC_*^{poly} of \mathcal{O}_M -modules placed in non-positive degrees

$$(1.36) \quad EC_k^{poly} = \begin{cases} \mathcal{O}_M, & \text{if } k = 0, \\ EJ_{-k-1}^{poly}, & \text{if } k < 0. \end{cases}$$

and the following \mathbb{R} -linear isomorphism of sheaves

$$(1.37) \quad \varrho : {}^E C_*^{poly} \rightarrow \ker \nabla^G \cap {}^E J_{-*}^{poly}$$

obtained by inverting the map (1.28).

Due to propositions 1.12 the action (1.31) and the differential \mathfrak{b} (1.35) commute with the Grothendieck connection ∇^G . Thus, the ∇^G -flat E -polyjets form a sheaf of DG submodule of $({}^E J_*^{poly}, \mathfrak{b}, {}^E S)$ over the sheaf of DGLAs $({}^E D_{poly}^*, \partial, [\cdot, \cdot]_G)$. Combining this observation with proposition 1.11 we conclude that the isomorphism (1.37) allows us to endow the sheaf (1.36) with a structure of a sheaf of DG modules over the sheaf of DGLAs ${}^E D_{poly}^*$. Namely,

Proposition 1.13. *The map*

$$(1.38) \quad {}^E R_\bullet : {}^E D_{poly}^k \otimes {}^E C_l^{poly} \rightarrow {}^E C_{k+l}^{poly}$$

given by the formula

$$(1.39) \quad {}^E R_P(a) = \chi^E S_P(\varrho(a)), \quad P \in \Gamma(U, {}^E D_{poly}^k), \quad a \in \Gamma(U, {}^E C_l^{poly})$$

and the differential

$$(1.40) \quad \mathfrak{b}(a) = \chi^E S_{1 \otimes 1}(\varrho(a)) : {}^E C_*^{poly} \rightarrow {}^E C_{*+1}^{poly}$$

turn ${}^E C_*^{poly}$ (1.36) into a sheaf of DG modules over the sheaf of DGLAs ${}^E D_{poly}^*$. \square

Remark 1. Since the map ϱ is NOT \mathcal{O}_M -linear the DGLA module structure (1.39), (1.40) on ${}^E C_*^{poly}$ is only \mathbb{R} -linear unlike the DGLA module structure (1.31) (1.35) on the sheaf ${}^E J_*^{poly}$.

Remark 2. It is not hard to see that in the case $E = TM$ the global sections of the sheaf ${}^E C_*^{poly}$ give the jet version [35] of the homological Hochschild complex of the algebra \mathcal{O}_M of functions on M .

The second remark motivates the following definition:

Definition 1.14. *We refer to the sheaf ${}^E C_*^{poly}$ of DG modules over the sheaf of DGLAs ${}^E D_{poly}^*$ of E -polydifferential operators as the sheaf of the Hochschild E -chains (or just E -chains for short).*

2. THE FORMALITY THEOREM FOR E -CHAINS

2.1. Hochschild-Kostant-Rosenberg. The cohomology of the complexes ${}^E D_{poly}^*$ and ${}^E C_*^{poly}$ are described by Hochschild-Kostant-Rosenberg type theorems. The original version of this theorem [19] says that the module of Hochschild homology of a smooth affine algebra is isomorphic to the module of exterior forms of the corresponding affine variety. In [4] A. Connes proved an analogous statement for the algebra of smooth functions on any compact real manifold, and in [34], N. Teleman was able to get rid of the assumption of compactness. The similar question about Hochschild cohomology turns out to be tractable if we replace the Hochschild cochains by polydifferential operators. We believe that the cohomology of this complex of polydifferential operators was originally computed by J. Vey [36]. All these computations correspond to the case when $E = TM$. In our general case we have the following proposition:

Proposition 2.1. *The natural maps*

$$(2.1) \quad \begin{aligned} \mathcal{V} : ({}^E T_{poly}^*, 0) &\longrightarrow ({}^E D_{poly}^*, \partial) \\ v_0 \wedge \cdots \wedge v_k &\longmapsto \frac{1}{(k+1)!} \sum_{\sigma \in S_{k+1}} \epsilon(\sigma) v_{\sigma_0} \otimes \cdots \otimes v_{\sigma_k} \end{aligned}$$

and

$$(2.2) \quad \begin{aligned} \mathfrak{C} : ({}^E C_*^{poly}, \mathfrak{b}) &\longrightarrow ({}^E A_*, 0) \\ a &\longmapsto (v \mapsto a \circ \mathcal{V}(v)) \end{aligned}$$

are quasi-isomorphisms of (sheaves of) complexes.

Remark. Recall that ${}^E A_*$ is the sheaf (1.7) of E -forms with reversed grading.

Proof. It is proved in [1] (Theorem 1.2) that \mathcal{V} is a quasi-isomorphism of cochain complexes. By (\mathcal{O}_M) -duality we obtain a quasi-isomorphism $({}^E C_*^{poly}, \mathfrak{b}) \rightarrow ({}^E A_*, 0)$, where $ba := a \circ \partial$. Let us show that $\mathfrak{b} = \mathfrak{b}$: let $a \in {}^E C_k^{poly}$ and $P \in {}^E D_{poly}^{k-1}$, then

$$\begin{aligned} (\mathfrak{b}a)(P) &= \rho(a)((1 \otimes P) \bullet m_0 + (-1)^{k-1}(1 \otimes P \otimes 1)) \\ &= \rho(a)(1 \otimes 1 \otimes P - 1 \otimes (P \bullet m_0) + (-1)^{k-1}(1 \otimes P \otimes 1)) \\ &= a(1 \otimes P - P \bullet m_0 + (-1)^{k-1}P \otimes 1) = a(\partial P). \end{aligned}$$

The proposition is proved. \square

2.2. The formality of the DGLA module of E -chains. Unfortunately, the maps (2.1) and (2.2) respect neither the Lie brackets nor the actions. This defect can be cured using the notion of Lie algebras and their modules *up to homotopy* (see [18] for a detailed discussion of the general theory and its applications, and [10, section 2] for a quick review of the notions and results we need). The main result of this paper is the following theorem:

Theorem 2.2. *For any C^∞ Lie algebroid (E, M, ρ) one can construct a commutative diagram of sheaves of DGLAs and DGLA modules over M*

$$(2.3) \quad \begin{array}{ccccccc} {}^E T_{poly}^* & \xrightarrow{\quad} & \mathcal{L}_1 & \xrightarrow{\quad} & \mathcal{L}_2 & \leftarrow \prec & {}^E D_{poly}^* \\ \downarrow \text{mod} & & \downarrow \text{mod} & & \downarrow \text{mod} & & \downarrow \text{mod} \\ {}^E A_* & \xrightarrow{\quad} & \mathcal{M}_1 & \leftarrow \prec \prec & \mathcal{M}_2 & \leftarrow \prec \prec & {}^E C_*^{poly} \end{array}$$

in which the horizontal arrows in the upper row are L_∞ -quasi-isomorphisms of sheaves of DGLAs and the horizontal arrows in the lower row are L_∞ -quasi-isomorphisms of L_∞ -modules. The terms $(\mathcal{L}_1, \mathcal{L}_2, \mathcal{M}_1, \mathcal{M}_2)$ and the quasi-isomorphisms of the diagram (2.3) are functorial for isomorphisms of pairs (E, ∂^E) , where E is a C^∞ Lie algebroid and ∂^E is a torsion free E -connection on E .

The proof of this theorem occupies the next two sections.

We would like to mention that the functoriality of the chain of quasi-isomorphisms (2.3) between the pair of sheaves of DGLA modules implies the following interesting results

Corollary 2.3. *Let (E, M, ρ) be a C^∞ Lie algebroid equipped with a smooth action of a group G . If one can construct a G -invariant connection ∂^E on E then there*

exists a chain of G -equivariant quasi-isomorphisms between the sheaves of DGLA modules $({}^E T_{poly}^*, {}^E A_*)$ and $({}^E D_{poly}^*, {}^E C_*^{poly})$. \square

In particular,

Corollary 2.4. *If (E, M, ρ) is a C^∞ Lie algebroid equipped with a smooth action of a finite or compact group G then the DGLA modules $(\Gamma(M, {}^E T_{poly}^*)^G, \Gamma(M, {}^E A_*)^G)$ and $(\Gamma(M, {}^E D_{poly}^*)^G, \Gamma(M, {}^E C_*^{poly})^G)$ are quasi-isomorphic.* \square

Example. Let us consider the case when the base manifold M shrinks to a point. Then the Lie algebroid E is a finite dimensional real Lie algebra \mathfrak{g} and the diagram of sheaves (2.3) becomes a diagram of (DG) Lie algebras and their modules. These DG Lie algebras and their modules can be described in geometric terms using a real Lie group G whose Lie algebra is \mathfrak{g} . Indeed, ${}^E D_{poly}^*$ can be identified with the DGLA of the left invariant polydifferential operators on G , ${}^E C_*^{poly}$ is the ${}^E D_{poly}^*$ -module of left invariant polyjets on G . Similarly, the sheaves ${}^E T_{poly}^*$ and ${}^E A_*$ can be identified with the graded Lie algebra of left invariant polyvector fields on G and the graded module of left invariant exterior forms on G , respectively. In this case, our result can be derived from corollary 4 in [11] (see section 5.3 in [11]).

Remark. It will appear clearly in the proof that all these results remain true for complex Lie algebroids. Namely, a complex Lie algebroid on a smooth real manifold M is a complex vector bundle of finite rank E whose sheaf of sections is a sheaf of (complex) Lie algebras with a $\mathcal{O}_M^{\mathbb{C}}$ -linear morphism of sheaves of Lie algebras $\rho : E \rightarrow T_{\mathbb{C}}M$ satisfying the same condition described in formula (1.1).

2.3. Formality theorems for the Hochschild complexes of $\mathbb{R}[[y^1, \dots, y^d]]$.

In order to prove theorem 2.2 we construct the Fedosov resolutions of the sheaves of DGLAs ${}^E T_{poly}^*$ and ${}^E D_{poly}^*$ and of the sheaves of DGLA modules ${}^E A_*$ and ${}^E C_*^{poly}$. These resolutions allow us to reduce the problem to the case of the tangent Lie algebroid $T\mathbb{R}^d \rightarrow \mathbb{R}^d$. For the latter case the desired result follows from the combination of Kontsevich's [20] and Shoikhet's [29] formality theorems.

First, we recall the required version of Kontsevich's formality theorem. Let $M = \mathbb{R}_{formal}^d$ be the formal completion of \mathbb{R}^d at the origin. In other words we set $\mathcal{O}_M = \mathbb{R}[[y^1, \dots, y^d]]$ and $E = \text{Der}(\mathcal{O}_M)$. Let us denote by $T_{poly}^*(\mathbb{R}_{formal}^d)$ and $D_{poly}^*(\mathbb{R}_{formal}^d)$ the DGLA of polyvector fields and polydifferential operators on \mathbb{R}_{formal}^d , respectively, then

Theorem 2.5 (Kontsevich, [20]). *There exists an L_∞ -quasi-isomorphism \mathcal{K} from $T_{poly}^*(\mathbb{R}_{formal}^d)$ to $D_{poly}^*(\mathbb{R}_{formal}^d)$ such that*

- (1) *The first structure map $\mathcal{K}^{[1]}$ is Vey's quasi-isomorphism (2.1) of complexes \mathcal{V} .*
- (2) *\mathcal{K} is $GL_d(\mathbb{R})$ -equivariant.*
- (3) *If $n > 1$ then for any vector fields $v_1, \dots, v_n \in T_{poly}^0(\mathbb{R}_{formal}^d)$*

$$\mathcal{K}^{[n]}(v_1, \dots, v_n) = 0$$

- (4) If $n > 1$ then for any vector field $v \in T_{poly}^0(\mathbb{R}_{formal}^d)$ linear in the coordinates y^i and any polyvector fields $\chi_2, \dots, \chi_n \in T_{poly}^*(\mathbb{R}_{formal}^d)$

$$\mathcal{K}^{[n]}(v, \chi_2, \dots, \chi_n) = 0.$$

We denote by

$$A^*(\mathbb{R}_{formal}^d) = \mathbb{R}[[y^1, \dots, y^d]] \otimes \bigwedge(\mathbb{R}^d)$$

the complex of exterior forms on \mathbb{R}_{formal}^d with the vanishing differential and by

$$J_*^{poly}(\mathbb{R}_{formal}^d) = \mathbb{R}[[y^1, \dots, y^d]]^{\hat{\otimes} (*+1)}$$

the complex of Hochschild chains of $\mathbb{R}[[y^1, \dots, y^d]]$, where the notation $\hat{\otimes}$ stands for the tensor product completed in the adic topology on $\mathbb{R}[[y^1, \dots, y^d]]$.

Using the Lie derivative (1.6) of exterior forms by a polyvector field, we can regard $A^*(\mathbb{R}_{formal}^d)$ as a graded module over the graded Lie algebra $T_{poly}^*(\mathbb{R}_{formal}^d)$. Furthermore, the action of Hochschild cochains on Hochschild chains (see formula (3.4) in [10]) allows us to regard $J_*^{poly}(\mathbb{R}_{formal}^d)$ as a DG modules over the DGLA $D_{poly}^*(\mathbb{R}_{formal}^d)$. Finally, using Kontsevich's quasi-isomorphism \mathcal{K} we get an L_∞ -module structure on $J_*^{poly}(\mathbb{R}_{formal}^d)$ over $T_{poly}^*(\mathbb{R}_{formal}^d)$. For this L_∞ -module, we have the following theorem:

Theorem 2.6 (Shoikhet, [29]). *There exists a quasi-isomorphism \mathcal{S} of L_∞ -modules over $T_{poly}^*(\mathbb{R}_{formal}^d)$ from $J_*^{poly}(\mathbb{R}_{formal}^d)$ to $A^*(\mathbb{R}_{formal}^d)$ such that*

- (1) *The first structure map $\mathcal{S}^{[1]}$ is the quasi-isomorphism of Connes (2.2).*
- (2) *The structure maps of \mathcal{S} are $GL_d(\mathbb{R})$ -equivariant.*
- (3) *If $n > 1$ then for any vector field $v \in T_{poly}^0(\mathbb{R}_{formal}^d)$ linear in the coordinates, any polyvector fields $\chi_2, \dots, \chi_n \in T_{poly}^*(\mathbb{R}_{formal}^d)$ and any chain $j \in J_*^{poly}(\mathbb{R}_{formal}^d)$*

$$\mathcal{S}^{[n]}(v, \chi_2, \dots, \chi_n; j) = 0$$

Remark 1. The third assertion of the above theorem is proved in [10] (see theorem 3).

Remark 2. Hopefully, one can prove the assertions of theorem 2.6 along the lines of Tamarkin and Tsygan [31, 32, 33].

3. THE FEDOSOV RESOLUTIONS

Let, as above, $E \rightarrow M$ be a C^∞ Lie algebroid with bracket $[\cdot, \cdot]$ on sections and the anchor ρ . Following [10] we introduce the formally completed symmetric algebra bundle $\hat{S}(E^\vee)$ of the dual bundle E^\vee and bundles \mathcal{T} , \mathcal{D} , \mathcal{A} , \mathcal{J} naturally associated to $\hat{S}(E^\vee)$. They all are pro- and/or ind-finite dimensional vector bundles.

- $\hat{S}(E^\vee)$ is the formally completed symmetric algebra bundle of the bundle E^\vee . Local sections are given by formal power series

$$\sum_{l=0}^{\infty} s_{i_1 \dots i_l}(x) y^{i_1} \dots y^{i_l}$$

where y^i are coordinates on the fibers of E and $s_{i_1 \dots i_l}$ are components of a symmetric covariant E -tensor.

- $\mathcal{T}^* := \hat{S}(E^\vee) \otimes \wedge^{*+1} E$ is the graded bundle of formal fiberwise polyvector fields on E . Local homogeneous sections of degree k are of the form

$$(3.1) \quad \sum_{l=0}^{\infty} v_{i_1 \dots i_l}^{j_0 \dots j_k}(x) y^{i_1} \dots y^{i_l} \frac{\partial}{\partial y^{j_0}} \wedge \dots \wedge \frac{\partial}{\partial y^{j_k}},$$

where $v_{i_1 \dots i_l}^{j_0 \dots j_k}$ are components of an E -tensor with symmetric covariant part (indices i_1, \dots, i_l) and antisymmetric contravariant part (indices j_0, \dots, j_k).

- $\mathcal{D}^* := \hat{S}(E^\vee) \otimes T^{*+1}(SE)$ is the graded bundle of formal fiberwise polydifferential operators on E with the shifted grading. A local homogeneous section of degree k looks as follow

$$(3.2) \quad \sum_{l=0}^{\infty} P_{i_1 \dots i_l}^{\alpha_0 \dots \alpha_k}(x) y^{i_1} \dots y^{i_l} \frac{\partial^{|\alpha_0|}}{\partial y^{\alpha_0}} \otimes \dots \otimes \frac{\partial^{|\alpha_k|}}{\partial y^{\alpha_k}},$$

where α_s are multi-indices, $P_{i_1 \dots i_l}^{\alpha_0 \dots \alpha_k}$ are components of an E -tensor with the obvious symmetry of the corresponding indices, and

$$\frac{\partial^{|\alpha_s|}}{\partial y^{\alpha_s}} = \frac{\partial}{\partial y^{j_1}} \dots \frac{\partial}{\partial y^{j_{|\alpha_s|}}}$$

for $\alpha_s = (j_1 \dots j_{|\alpha_s|})$.

- $\mathcal{A}_* := \hat{S}(E^\vee) \otimes \wedge^{-*}(E^\vee)$ is the graded bundle of formal fiberwise differential forms on E with the reversed grading. Any local homogeneous section of degree $-k$ can be written as

$$(3.3) \quad \sum_{l=0}^{\infty} \omega_{i_1 \dots i_l, j_1 \dots j_k}(x) y^{i_1} \dots y^{i_l} dy^{j_1} \wedge \dots \wedge dy^{j_k},$$

where $\omega_{i_1 \dots i_l, j_1 \dots j_k}$ are components of a covariant E -tensor symmetric in indices i_1, \dots, i_l and antisymmetric in indices j_1, \dots, j_k .

- \mathcal{J}_* is the bundle of Hochschild chains of $\hat{S}(E^\vee)$ over \mathcal{O}_M .

$$(3.4) \quad \mathcal{J} = \bigoplus_{k \geq 0} \mathcal{J}_k, \quad \mathcal{J}_k := (\hat{S}E^\vee)^{\hat{\otimes}_{\mathcal{O}_M} (k+1)},$$

where $\hat{\otimes}$ stands for the tensor product completed in the adic topology. Local sections of homogeneous degree k are formal power series

$$(3.5) \quad \sum_{\alpha_0, \dots, \alpha_k} a_{\alpha_0, \dots, \alpha_k}(x) y_0^{\alpha_0} y_1^{\alpha_1} \dots y_k^{\alpha_k}$$

in $k+1$ copies y_0, \dots, y_k of coordinates on the fibers of E . Here α_s are multi-indices, $a_{\alpha_0, \dots, \alpha_k}$ are components of a tensor with an obvious symmetry in the corresponding indices, and

$$y_m^{\alpha_m} = y_m^{j_1} \dots y_m^{j_{|\alpha_m|}}$$

for $\alpha_m = (j_1 \dots j_{|\alpha_m|})$.

For our purposes, we consider E -differential forms with values in the sheaves $\hat{S}(E^\vee)$, \mathcal{T} , \mathcal{D} , \mathcal{A} , \mathcal{J} . Below we list these sheaves of E -forms together with the algebraic structures they carry.³

³For any bundle \mathcal{B} we will denote ${}^E\Omega(\mathcal{B})$ the bundle ${}^E\Omega \otimes \mathcal{B}$ of E -forms with values in \mathcal{B} , and ${}^E\Omega(U, \mathcal{B})$ the space of sections over an open subset $U \subset M$ (instead of $\Gamma(U, {}^E\Omega \otimes \mathcal{B})$).

- ${}^E\Omega(\hat{S}(E^\vee))$ is a bundle of graded commutative algebras with grading given by the exterior degree of E -forms. ${}^E\Omega(\hat{S}(E^\vee))$ is also filtered by the degree of monomials in fiber coordinates y^i .
- ${}^E\Omega(\mathcal{T})$ is a sheaf of graded Lie algebras and ${}^E\Omega(\mathcal{A})$ is a sheaf of graded modules over ${}^E\Omega(\mathcal{T})$. These structures are induced by those of $T_{poly}^*(\mathbb{R}_{formal}^d)$ and $A^*(\mathbb{R}_{formal}^d)$, respectively and the grading is given by the sum of the exterior degree and the degree of a polyvector (resp. a form). $[\cdot, \cdot]_{SN}$ will denote the Lie bracket between sections of the sheaf ${}^E\Omega(\mathcal{T})$ and L_u (the Lie derivative) will denote the action of a fiberwise polyvector $u \in {}^E\Omega(\mathcal{T})$ on the sections of ${}^E\Omega(\mathcal{A})$. ${}^E\Omega(\mathcal{T})$ is also a sheaf of graded commutative algebras. The multiplication of sections in ${}^E\Omega(\mathcal{T})$ is given by the exterior product in the space $T_{poly}^*(\mathbb{R}_{formal}^d)$ of polyvector fields on \mathbb{R}_{formal}^d . The Lie bracket and the product in ${}^E\Omega(\mathcal{T})$ turn ${}^E\Omega(\mathcal{T})$ into a sheaf of Gerstenhaber algebras⁴.
- ${}^E\Omega(\mathcal{D})$ is a sheaf of DGLAs and ${}^E\Omega(\mathcal{J})$ is a sheaf of DGLA modules over ${}^E\Omega(\mathcal{D})$. These structures are induced by those of $D_{poly}^*(\mathbb{R}_{formal}^d)$ and $J_{poly}^*(\mathbb{R}_{formal}^d)$, respectively and the grading is given by the sum of the exterior degree and the degree of a (co)chain. We denote by ∂ and $[\cdot, \cdot]_G$ respectively the differential and the Lie bracket on ${}^E\Omega(\mathcal{D})$, \mathfrak{b} will stand for the differential on ${}^E\Omega(\mathcal{J})$ and \mathcal{R}_P will denote the action of $P \in {}^E\Omega(\mathcal{D})$ on the sections of ${}^E\Omega(\mathcal{J})$. ${}^E\Omega(\mathcal{D})$ is also a sheaf of DGAs. The multiplication of sections is induced by the cup product in the space $D_{poly}^*(\mathbb{R}_{formal}^d)$ of polydifferential operators on \mathbb{R}_{formal}^d .

Remark. Notice that \mathcal{A} is a sheaf of exterior forms with values in $\hat{S}(E^\vee)$. However, we would like to distinguish \mathcal{A} from ${}^E\Omega(\hat{S}(E^\vee))$. For this purpose we use two copies of a local basis of exterior forms. Those are $\{dy^i\}$ and $\{\xi^i\}$ for \mathcal{A} and ${}^E\Omega(\hat{S}(E^\vee))$, respectively.

The following proposition shows that we have a distinguished sheaf of graded Lie algebras which acts on the sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$.

Proposition 3.1. *The sheaf ${}^E\Omega(\mathcal{T}^0)$ of E -forms with values in fiberwise vector fields is a sheaf of graded Lie algebras. The sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$ are sheaves of modules over ${}^E\Omega(\mathcal{T}^0)$ and the action of sections in ${}^E\Omega(\mathcal{T}^0)$ is compatible with the DG algebraic structures on ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$.*

Proof. Since the Schouten-Nijenhuis bracket (1.3), (1.4) has degree zero ${}^E\Omega(\mathcal{T}^0) \subset {}^E\Omega(\mathcal{T}) \subset {}^E\Omega(\mathcal{D})$ is a subsheaf of graded Lie algebras. While the action of ${}^E\Omega(\mathcal{T}^0)$ on the sections of ${}^E\Omega(\hat{S}(E^\vee))$ is obvious, the action on ${}^E\Omega(\mathcal{A})$ is given by the Lie derivative, the action on ${}^E\Omega(\mathcal{T})$ is the adjoint action corresponding to the Schouten-Nijenhuis bracket, the action on ${}^E\Omega(\mathcal{D})$ is given by the Gerstenhaber bracket and the action on ${}^E\Omega(\mathcal{J})$ is induced by the action of Hochschild cochains on Hochschild chains (see formula 3.4 in paper [10]). The compatibility of the action with the corresponding DGLA and DGLA module structures follows from the construction. The compatibility of the action with the product in ${}^E\Omega(\mathcal{T})$ follows from the axioms

⁴The definition of the Gerstenhaber algebra can be found in section 4.1 of the second part of [7] or in the original paper [16].

of the Gerstenhaber algebra [16] and the compatibility with the product in ${}^E\Omega(\mathcal{D})$ can be verified by a straightforward computation. \square

Due to the above proposition the following 2-nilpotent derivation

$$(3.6) \quad \delta := \xi^i \frac{\partial}{\partial y^i} : {}^E\Omega^*(\hat{S}(E^\vee)) \rightarrow {}^E\Omega^{*+1}(\hat{S}(E^\vee))$$

of the sheaf of algebras ${}^E\Omega(\hat{S}(E^\vee))$ obviously extends to 2-nilpotent differentials on ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$, ${}^E\Omega(\mathcal{A})$ and ${}^E\Omega(\mathcal{J})$. Furthermore, it follows from proposition 3.1 that δ is compatible with the DG algebraic structures on ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$.

Note that

$$(3.7) \quad \ker \delta \cap \hat{S}(E^\vee) \cong \mathcal{O}_M, \quad \ker \delta \cap \mathcal{A}_* \cong {}^E\mathcal{A}_*$$

as sheaves of (graded) commutative algebras over \mathcal{O}_M . Similarly, $\ker \delta \cap \mathcal{T}$, (resp. $\ker \delta \cap \mathcal{D}$) is a sheaf of fiberwise polyvector fields (3.1) (resp. fiberwise polydifferential operators (3.2)) whose components do not depend on the fiber coordinates y^i . In other words,

$$(3.8) \quad \ker \delta \cap \mathcal{T}^* \cong \wedge^{*+1}(E)$$

as sheaves of graded commutative algebras and

$$(3.9) \quad \ker \delta \cap \mathcal{D}^* \cong \otimes^{*+1}(S(E)),$$

as sheaves of DGAs over \mathcal{O}_M .

In fact, one can prove a more stronger statement:

Proposition 3.2. *For \mathcal{B} being either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} or \mathcal{D}*

$$H^{\geq 1}({}^E\Omega(\mathcal{B}), \delta) = 0.$$

Furthermore,

$$(3.10) \quad \begin{aligned} H^0({}^E\Omega(\hat{S}(E^\vee)), \delta) &\cong \mathcal{O}_M, \\ H^0({}^E\Omega(\mathcal{A}_*), \delta) &\cong {}^E\mathcal{A}_*, \\ H^0({}^E\Omega(\mathcal{T}^*), \delta) &\cong \wedge^{*+1}(E) \end{aligned}$$

as sheaves of (graded) commutative algebras and

$$(3.11) \quad H^0({}^E\Omega(\mathcal{D}^*), \delta) \cong \otimes^{*+1}(S(E))$$

as sheaves of DGAs over \mathcal{O}_M .

Proof. Due to equations (3.7), (3.8), and (3.9) the proposition will follow immediately if we construct an operator

$$(3.12) \quad \kappa : {}^E\Omega^*(\mathcal{B}) \rightarrow {}^E\Omega^{*-1}(\mathcal{B})$$

such that for any section u of ${}^E\Omega(\mathcal{B})$

$$(3.13) \quad u = \delta\kappa(u) + \kappa\delta(u) + \mathcal{H}(u),$$

where

$$(3.14) \quad \mathcal{H}(u) = u \Big|_{y^i = \xi^i = 0}.$$

First, we define this operator on the sheaf ${}^E\Omega(\hat{S}(E^\vee))$

$$(3.15) \quad \kappa(a) = y^k \frac{\bar{\partial}}{\partial \xi^k} \int_0^1 a(x, ty, t\xi) \frac{dt}{t}, \quad a \in {}^E\Omega^{>0}(U, \hat{S}(E^\vee)), \quad \kappa|_{\hat{S}(E^\vee)} = 0,$$

where the arrow over ∂ denotes the left derivative with respect to the anti-commuting variable ξ^i .

Next, we extend κ to sections of the sheaves ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$ in the componentwise manner. A direct computation shows that equation (3.13) holds and the proposition follows. \square

Since our Lie algebroid E is a smooth bundle over M , it admits a global torsion free connection ∂^{E^5} . Using this connection we define the following derivation of the DG sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$:

$$(3.16) \quad \nabla = {}^E d + \Gamma \cdot : {}^E\Omega^*(\mathcal{B}) \rightarrow {}^E\Omega^{*+1}(\mathcal{B}), \quad \Gamma = -\xi^i \Gamma_{ij}^k y^j \frac{\partial}{\partial y^k},$$

where \mathcal{B} is either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , \mathcal{D} , or \mathcal{J} , $\Gamma_{ij}^k(x)$ are Christoffel's symbols of the connection ∂^E and $\Gamma \cdot$ denotes the action of Γ on the sections of the sheaves ${}^E\Omega(\mathcal{B})$ (see proposition 3.1). It is not hard to see that ∇ (3.16) is compatible with the DG algebraic structures on ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$. Furthermore, the torsion freeness of the connection ∂^E implies that

$$(3.17) \quad \nabla \delta + \delta \nabla = 0.$$

The standard curvature E -tensor $(R_{ij})^l_k(x)$ of the connection ∂^E provides us with the following fiberwise vector field:

$$(3.18) \quad R = -\frac{1}{2} \xi^i \xi^j (R_{ij})^l_k(x) y^k \frac{\partial}{\partial y^l} \in {}^E\Omega^2(M, \mathcal{T}^0).$$

A direct computation shows that for \mathcal{B} being any of the sheaves ${}^E S$, \mathcal{A} , \mathcal{T} , \mathcal{D} , or \mathcal{J} , we have

$$(3.19) \quad \nabla^2 = R \cdot : {}^E\Omega^*(\mathcal{B}) \rightarrow {}^E\Omega^{*+2}(\mathcal{B}),$$

where $R \cdot$ denotes the action of the vector field R in the sense of proposition 3.1.

Although ∇ is not flat the following theorem shows that the combination $\nabla - \delta$ can be extended to a flat connection on the sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{D})$, and ${}^E\Omega(\mathcal{J})$.

Theorem 3.3. *Let \mathcal{B} be either of the sheaves ${}^E S$, \mathcal{A} , \mathcal{T} , \mathcal{D} , or \mathcal{J} . There exists a global section*

$$(3.20) \quad A = \sum_{s=2}^{\infty} \xi^k A_{k, i_1 \dots i_s}^j(x) y^{i_1} \dots y^{i_s} \frac{\partial}{\partial y^j}$$

of the sheaf ${}^E\Omega^1(\mathcal{T}^0)$ such that the derivation

$$(3.21) \quad D := \nabla - \delta + A \cdot : {}^E\Omega^*(\mathcal{B}) \rightarrow {}^E\Omega^{*+1}(\mathcal{B})$$

is 2-nilpotent

$$D^2 = 0,$$

and (3.21) is compatible with the DG algebraic structure on ${}^E\Omega(\mathcal{B})$.

⁵Recall that by the word ‘‘connection’’ we always mean an E -connection (1.14).

Proof. The proof goes essentially along the lines of [9, theorem 2].

Thanks to equation (3.19) the condition $D^2 = 0$ is equivalent to the equation

$$(3.22) \quad R + \nabla A - \delta A + \frac{1}{2}[A, A]_{SN} = 0.$$

We claim that a solution of (3.22) can be obtained by iterations of the following equation

$$(3.23) \quad A = \kappa R + \kappa(\nabla A + \frac{1}{2}[A, A]_{SN})$$

in degrees in the fiber coordinates y^i . Indeed, equation (3.13) implies that iterating (3.23) we get a solution of the equation

$$\kappa(R + \nabla A - \delta A + \frac{1}{2}[A, A]_{SN}) = 0.$$

We denote by C the left hand side of (3.22)

$$C = R + \nabla A - \delta A + \frac{1}{2}[A, A]_{SN},$$

and mention that due to Bianchi's identities $\nabla R = \delta R = 0$

$$(3.24) \quad \nabla C - \delta C + [A, C] = 0.$$

Applying κ (3.15) to (3.24) and using the homotopy property (3.13) we get

$$C = \kappa(\nabla C + [A, C]).$$

The latter equation has the unique vanishing solution since the operator κ (3.15) raises the degree in the fiber coordinates y^i .

Proposition 3.1 implies that the differential (3.21) is compatible with the DG algebraic structures on ${}^E\Omega(\mathcal{B})$. Thus, the theorem is proved. \square

In what follows we refer to the differential D (3.21) as *the Fedosov differential*.

The following theorem describes the cohomology of the Fedosov differential D for the sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, and ${}^E\Omega(\mathcal{D})$

Theorem 3.4. *For \mathbf{B} being either of the sheaves ${}^E\Omega(\hat{S}(E^\vee))$, ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, or ${}^E\Omega(\mathcal{D})$*

$$(3.25) \quad H^{\geq 1}(\mathbf{B}, D) = 0.$$

Furthermore,

$$(3.26) \quad \begin{aligned} H^0({}^E\Omega(\hat{S}(E^\vee)), D) &\cong \mathcal{O}_M, \\ H^0({}^E\Omega(\mathcal{A}_*), D) &\cong {}^E\mathcal{A}_*, \\ H^0({}^E\Omega(\mathcal{T}^*), D) &\cong \ker \delta \cap \mathcal{T}^*, \end{aligned}$$

as sheaves of graded commutative algebras

$$(3.27) \quad H^0({}^E\Omega(\mathcal{D}^*), D) \cong \ker \delta \cap \mathcal{D}^*$$

as sheaves of DGAs over \mathbb{R} .

Proof. The first statement follows easily from the spectral sequence argument. Indeed, using the fiber coordinates y^i we introduce the decreasing filtration

$$\dots \subset F^{p+1}\mathbf{B} \subset F^p\mathbf{B} \subset F^{p-1}\mathbf{B} \subset \dots \subset F^0\mathbf{B} = \mathbf{B},$$

where the components of the sections of the sheaf $F^p\mathbf{B}$ have degree in $y^i \geq p$.

Since $D(F^p\mathbf{B}) \subset F^{p-1}\mathbf{B}$ the corresponding spectral sequence starts with

$$E_{-1}^{p,q} = F^p\mathbf{B}^{p+q}.$$

It is easy to see that

$$d_{-1} = \delta.$$

Thus using proposition 3.2 we conclude that for any p, q satisfying the condition $p + q > 0$

$$E_0^{p,q} = E_1^{p,q} = \dots = E_\infty^{p,q} = 0$$

and the first statement (3.25) follows.

Let \mathcal{B} denote either of the bundles $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , or \mathcal{D} . We claim that iterating the equation

$$(3.28) \quad \lambda(u) = u + \kappa(\nabla\lambda(u) + A \cdot \lambda(u)), \quad u \in \Gamma(U, \mathcal{B} \cap \ker \delta)$$

we get a map of sheaves of graded vector spaces

$$(3.29) \quad \lambda : \mathcal{B} \cap \ker \delta \rightarrow \mathcal{B} \cap \ker D.$$

Here $A \cdot$ denotes the action of the fiberwise vector field A , defined in proposition 3.1. Indeed, let u be a section of \mathcal{B} . Then, due to formula (3.13) $\lambda(u)$ satisfies the following equation

$$(3.30) \quad \kappa(D(\lambda(u))) = 0.$$

Let us denote $D\lambda(u)$ by Y

$$Y = D\lambda(u).$$

The equation $D^2 = 0$ implies that

$$DY = 0$$

which is equivalent to

$$(3.31) \quad \delta Y = \nabla Y + A \cdot Y$$

Applying (3.13) to Y and using equations (3.30), (3.31) we get

$$Y = \kappa(\nabla Y + A \cdot Y).$$

The latter equation has the unique vanishing solution since the operator κ (3.15) raises the degree in the fiber coordinates y^i .

The map (3.29) is obviously injective. To prove that the map is surjective we notice that \mathcal{H}

$$\mathcal{H} : \mathcal{B} \rightarrow \mathcal{B} \cap \ker \delta$$

is a left inverse of the map (3.29). Thus it suffices to prove that if $a \in \Gamma(U, \mathcal{B} \cap \ker D)$ and

$$(3.32) \quad \mathcal{H}a = 0$$

then a vanishes.

The condition $a \in \ker D$ is equivalent to the equation

$$\delta a = \nabla a + A \cdot a.$$

Hence, applying (3.13) to a and using (3.32) we get

$$a = \kappa(\nabla a + A \cdot a).$$

The latter equation has the unique vanishing solution since the operator κ (3.15) raises the degree in the fiber coordinates y^i . Thus, the map (3.29) is bijective and the map \mathcal{H}

$$(3.33) \quad \mathcal{H} : \mathcal{B} \cap \ker D \rightarrow \mathcal{B} \cap \ker \delta$$

is the inverse of (3.29).

It remains to prove that the map (3.29) is compatible with the multiplication of the sections of the sheaf \mathcal{B} , where \mathcal{B} is either $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , or \mathcal{D} . The latter follows immediately from the fact that the inverse map \mathcal{H}

$$(3.34) \quad \mathcal{H} : \mathcal{B} \rightarrow \mathcal{B} \cap \ker \delta$$

respects the corresponding algebra structures on $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , and the DGA structure on \mathcal{D} . \square

Let us now mention that since the Fedosov differential (3.21) is compatible with the graded algebraic structures on the sheaves ${}^E\Omega(\mathcal{T})$ and ${}^E\Omega(\mathcal{A})$ we conclude that $H^*({}^E\Omega(\mathcal{T}), D)$ is a sheaf of graded Lie algebras and $H^*({}^E\Omega(\mathcal{A}), D)$ is a sheaf of graded modules over $H^*({}^E\Omega(\mathcal{T}), D)$. On the other hand the above theorem tells us that

$$H^*({}^E\Omega(\mathcal{A}), D) = {}^EA_*,$$

and

$$H^*({}^E\Omega(\mathcal{T}), D) = \mathcal{T}^* \cap \ker \delta,$$

Furthermore, the sheaf $\mathcal{T}^* \cap \ker \delta$ in the right hand side of the latter equation can be canonically identified with ${}^ET_{poly}^* = \wedge^{*+1} E$ as a sheaf of vector spaces.

Thus, it is natural to ask whether the graded algebraic structures on the sheaves $\mathcal{T}^* \cap \ker \delta$ and EA_* coincide with the ones given by Lie bracket (1.3) (1.4) and the Lie derivative (1.6). A positive answer to this question is given by the following proposition:

Proposition 3.5. *The composition*

$$(3.35) \quad \mathcal{H}' = \nu \circ \mathcal{H} : \mathcal{T}^* \cap \ker D \rightarrow {}^ET_{poly}^*$$

of the map

$$(3.36) \quad \mathcal{H} : \mathcal{T}^* \cap \ker D \rightarrow \mathcal{T}^* \cap \ker \delta$$

with the identification of the sheaves $\mathcal{T}^ \cap \ker \delta$ and ${}^ET_{poly}^* \cong \wedge^{*+1} E$*

$$(3.37) \quad \nu : \mathcal{T}^* \cap \ker \delta \xrightarrow{\cong} {}^ET_{poly}^*$$

induces an isomorphism of the sheaves of graded Lie algebras $H^({}^E\Omega(\mathcal{T}), D) \cong {}^ET_{poly}^*$. The map*

$$(3.38) \quad \mathcal{H} : \mathcal{A}_* \cap \ker D \rightarrow {}^EA_*$$

induces an isomorphism of the sheaves of graded modules $H^({}^E\Omega(\mathcal{A}), D) \cong {}^EA_*$ over the sheaf of graded Lie algebras $H^*({}^E\Omega(\mathcal{T}), D) \cong {}^ET_{poly}^*$.*

Proof. The first part of the proposition is proved in [1] (see proposition 2.4). To prove the second part, we first remark that the maps \mathcal{H} and ν are compatible with the cup products.

Next, we show that for any D -closed fiberwise differential form $\omega \in \Gamma(U, \mathcal{A})$ one has

$$\mathcal{H}(d^f \omega) = {}^E d \mathcal{H}(\omega),$$

where $d^f = dy^i \frac{\partial}{\partial y^i}$ is the fiberwise De Rham differential on \mathcal{A} . Since

$$\mathcal{H} : \mathcal{A}_* \rightarrow {}^E A_*$$

is a morphism of graded commutative algebras, it is sufficient to prove it for functions and 1-forms:

- *First case.* Let f be a function and

$$\omega = \lambda(f).$$

A direct computation shows that

$$\lambda(f) = f + y^i \rho(e_i) f \pmod{|y|^2}.$$

Therefore $d^f \omega = \rho(e_i) f dy^i \pmod{|y|}$, and hence, $\mathcal{H}(d^f \omega) = {}^E df$.

- *Second case.* Let $\alpha = \alpha_i(x) dy^i$ be a E -1-form and

$$\omega = \lambda(\alpha).$$

It is not hard to show that

$$\lambda(\alpha) = \alpha + y^i (\rho(e_i) \alpha_j - \Gamma_{ij}^k \alpha_k) dy^j \pmod{|y|^2}.$$

Therefore,

$$d^f \omega = (\rho(e_i) \alpha_j - \Gamma_{ij}^k \alpha_k) dy^i \wedge dy^j \pmod{|y|} = (\rho(e_i) \alpha_j - \frac{1}{2} c_{ij}^k \alpha_k) dy^i \wedge dy^j \pmod{|y|},$$

and hence,

$$\mathcal{H}(d^f \omega) = {}^E d \alpha.$$

To finish the proof we notice that for any fiberwise polyvector field $u \in \Gamma(U, \mathcal{T}^*)$ and any fiberwise differential form $\omega \in \Gamma(U, \mathcal{A})$, the equation

$$\mathcal{H}(\iota_u \omega) = \iota_{\mathcal{H}(u)} \circ \mathcal{H}(\omega)$$

is obviously satisfied. The latter implies that for any pair of D -closed sections $u \in \Gamma(U, \mathcal{T}^*)$, $\omega \in \Gamma(U, \mathcal{A}_*)$

$$\mathcal{H}(L_u \omega) = {}^E L_{\mathcal{H}(u)} \circ \mathcal{H}(\omega),$$

and the proposition follows. \square

Remark. Actually, we have proved a slightly stronger statement. Namely, we shown that the maps (3.38) and (3.35) induce an isomorphism of the sheaves of *calculi*.

$$(H^*({}^E \Omega(\mathcal{T}), D), H^*({}^E \Omega(\mathcal{A}), D)) \cong ({}^E T_{poly}^*, {}^E A_*).$$

The precise definition of the calculi can be found in section 4.3 of the second part of [7].

Let us now recall that \mathcal{T}^0 is a sheaf of Lie-Rinehart algebras [28] over the sheaf of algebras $\mathcal{T}^{-1} = \hat{S}(E^\vee)$, and \mathcal{D}^0 is the universal enveloping algebra⁶ of \mathcal{T}^0 . Therefore, the inverse $(\mathcal{H}')^{-1}$ of the map (3.35) induces the morphism

$$(3.39) \quad \mu : \mathcal{U}E \rightarrow \mathcal{D}^0$$

of the sheaves of Hopf algebroids with anchor, and for any $P \in \Gamma(U, \mathcal{U}E)$

$$(3.40) \quad D(\mu(P)) = 0.$$

We claim that

Proposition 3.6. *The map (3.39) gives the isomorphism*

$$(3.41) \quad \mu : \mathcal{U}E \rightarrow \mathcal{D}^0 \cap \ker D.$$

of the sheaves of Hopf algebroids with anchor.

Proof. Notice that $\mathcal{U}E$ and \mathcal{D}^0 are both filtered sheaves of algebras. The filtration on $\mathcal{U}E$ is defined in (1.12) and the filtration on \mathcal{D}^0 is given by the degree of differential operators.

Thanks to the results of [27] and [28] we have the PBW theorem for Lie algebroids. This theorem says that the associated graded module of the filtration (1.12) on $\mathcal{U}E$ is

$$Gr(\mathcal{U}E) = S(E)$$

the symmetric algebra of the bundle E .

Furthermore, it is not hard to see that the map μ is compatible with the filtrations on $\mathcal{U}E$ and \mathcal{D}^0 and due to theorem 3.4 and proposition 3.2 μ induces the isomorphism

$$S(E) \cong \mathcal{D}^0 \cap \ker D$$

of the associated graded sheaves of vector spaces. Therefore, the snake lemma argument implies that the map (3.41) is also an isomorphism onto the sheaf $\mathcal{D}^0 \cap \ker D$ of D -flat sections of \mathcal{D}^0 . \square

Let us recall that ${}^E D_{poly}^*$ (resp. \mathcal{D}^*) is the tensor algebras of $\mathcal{U}E$ over \mathcal{O}_M (resp. the tensor algebra of \mathcal{D}^0 over $\hat{S}(E^\vee)$). Using this fact we extend (3.39) to the morphism

$$(3.42) \quad \mu' : {}^E D_{poly}^* \rightarrow \mathcal{D}^*.$$

of sheaves of DGAs (over \mathbb{R}) by setting

$$\mu' \Big|_{{}^E D_{poly}^0} = \mu, \quad \mu' \Big|_{\mathcal{O}_M} = \lambda,$$

where λ is defined in (3.29).

Let us also observe that since the map (3.39) is a morphism of the sheaves of Hopf algebroids with anchor then the map (3.42) a morphism of the sheaves of DGLAs (over \mathbb{R}). Furthermore, theorem 3.4 implies that the sheaf of DGAs $\mathcal{D}^* \cap \ker D$ is generated by the sheaf $\mathcal{D}^0 \cap \ker D$ over the sheaf of commutative algebras $\hat{S}(E^\vee) \cap \ker D \cong \mathcal{O}_M$. Therefore using proposition 3.6 we get the following result:

⁶More precisely, \mathcal{D}^0 is the sheaf associated to the corresponding presheaf of universal enveloping algebras, like $\mathcal{U}E$ (1.11) for the sheaf of Lie-Rinehart algebras E over \mathcal{O}_M .

Proposition 3.7 (proposition 2.5, [1]). *The map (3.42) gives an isomorphism of the sheaves of DGLAs*

$$(3.43) \quad \mu' : {}^E D_{poly}^* \xrightarrow{\sim} \mathcal{D}^* \cap \ker D.$$

This map is also compatible with the DGA structures on the sheaves ${}^E D_{poly}^$ and $\mathcal{D}^* \cap \ker D$ by construction. \square*

Let us consider the map of sheaves of graded vector spaces

$$(3.44) \quad \gamma : \mathcal{J}_* \rightarrow {}^E J_*^{poly}, \quad \gamma(j)(P) = (\mu'(P))(j) \Big|_{y^i=0},$$

$$j \in \Gamma(U, \mathcal{J}_k), \quad P \in \Gamma(U, {}^E D_{poly}^k).$$

We claim that

Theorem 3.8. *For any $q \geq 1$*

$$(3.45) \quad H^q({}^E \Omega(\mathcal{J}), D) = 0,$$

and the map (3.44) gives an isomorphism of the sheaves of DG modules over the sheaf of DGLAs ${}^E D_{poly}^ \cong \mathcal{D}^* \cap \ker D$*

$$(3.46) \quad \gamma : \mathcal{J}_* \xrightarrow{\sim} {}^E J_*^{poly}.$$

This isomorphism sends the Fedosov connection (3.21) on \mathcal{J}^ to the Grothendieck connection (1.26) on ${}^E J_*^{poly}$.*

Proof. The first statement (3.45) follows easily from the spectral sequence argument. Indeed, using the zeroth collection of the fiber coordinates y_0^i (3.5) we introduce the decreasing filtration on the sheaf ${}^E \Omega(\mathcal{J})$

$$\dots \subset F^{p+1}({}^E \Omega(\mathcal{J})) \subset F^p({}^E \Omega(\mathcal{J})) \subset F^{p-1}({}^E \Omega(\mathcal{J})) \subset \dots \subset F^0({}^E \Omega(\mathcal{J})) = {}^E \Omega(\mathcal{J}),$$

where the components of the sections (3.5) of the sheaf $F^p({}^E \Omega(\mathcal{J}))$ have degree in $y_0^i \geq p$.

Since $D(F^p({}^E \Omega(\mathcal{J}))) \subset F^{p-1}({}^E \Omega(\mathcal{J}))$ the corresponding spectral sequence starts with

$$E_{-1}^{p,q} = F^p({}^E \Omega(\mathcal{J})^{p+q}).$$

Next, we observe that

$$d_{-1} = \xi^i \frac{\partial}{\partial y_0^i},$$

and hence, due to the Poincaré lemma for the formal disk we have

$$E_0^{p,q} = E_1^{p,q} = \dots = E_\infty^{p,q} = 0$$

whenever $p+q > 0$. Thus, the first statement (3.45) of the theorem follows.

Since (3.39) is a morphism of sheaves of Hopf algebroids with anchor

$$\mu'(P \bullet Q) = \mu'(P) \bullet \mu'(Q), \quad P, Q \in \Gamma(U, {}^E D_{poly}^*).$$

Furthermore, μ' is obviously compatible with cyclic permutations

$$t \mu'(P_0 \otimes P_1 \otimes \dots \otimes P_l) = \mu'(P_1 \otimes P_2 \otimes \dots \otimes P_l \otimes P_0), \quad P_i \in \Gamma(U, \mathcal{U}E).$$

Hence, for any $P \in \Gamma(U, {}^E D_{poly}^*)$ and any $a \in \Gamma(U, \mathcal{J}_*)$

$$(3.47) \quad {}^E S_P(\gamma(a)) = \gamma(\mathcal{R}_{\mu'(P)}(a)).$$

Since \mathcal{J}_* is dual to $\mathcal{D}^* \cap \ker \delta$ and $\mathcal{D}^* \cap \ker \delta \cong \mathcal{D}^* \cap \ker D \cong {}^E D_{poly}^*$ the map (3.46) is an isomorphism. It remains to prove that the map (3.46) sends the Fedosov

connection (3.21) to the Grothendieck connection (1.26). This statement is proved by the following line of equations:

$$\begin{aligned}
 \gamma(D_u j)(P) &= (\mu'(P))(D_u j) \Big|_{y^i=0} = (D_u[\mu'(P)(j)]) \Big|_{y^i=0} \\
 &= \rho(u)[\mu'(P)(j)] \Big|_{y^i=0} - (\iota_u \delta \bullet [\mu'(P)(j)]) \Big|_{y^i=0} \\
 &= \rho(u)[\mu'(P)(j)] \Big|_{y^i=0} - (\mu'(u) \bullet \mu'(P)(j)) \Big|_{y^i=0} \\
 &= \rho(u)[\mu'(P)(j)] \Big|_{y^i=0} - \mu'(u \bullet P)(j) \Big|_{y^i=0} \\
 &= \rho(u)(\gamma(j))(P) - (\gamma(j))(u \bullet P) = (\nabla_u^G \gamma(j))(P),
 \end{aligned}$$

where $u \in \Gamma(U, E)$, $j \in \Gamma(U, \mathcal{J}_k)$, $P \in \Gamma(U, {}^E D_{poly}^k)$, ι denotes the contraction of an E -vector field with E -differential forms, ρ is the anchor map, and u is viewed both as a section of E and an E -differential operator. \square

4. PROOF OF THE FORMALITY THEOREM FOR E -CHAINS AND ITS APPLICATIONS

4.1. **Proof of the theorem.** Let us denote

- $\lambda_A : {}^E A_* \rightarrow {}^E \Omega(\mathcal{A}_*)$ the map λ (3.29) defined in the proof of theorem 3.4 for $\mathcal{B} = {}^E \Omega(\mathcal{A}_*)$,
- $\lambda_T : {}^E T_{poly}^* \rightarrow {}^E \Omega(\mathcal{T})$, the inverse of the map \mathcal{H}' (3.35),
- $\lambda_D : {}^E D_{poly}^* \rightarrow {}^E \Omega(\mathcal{D})$, the map μ' (3.42) and
- $\lambda_C : {}^E C_*^{poly} \rightarrow {}^E \Omega(\mathcal{J})$, the composition $\gamma^{-1} \circ \varrho$ of the inverse of the map γ (3.44) with the map ϱ (1.37).

The results of the previous section can be represented in the form of the following commutative diagrams of sheaves of DGLAs, their modules, and morphisms

$$\begin{array}{ccc}
 ({}^E T_{poly}^*, 0, [,]_{SN}) & \xrightarrow{\lambda_T} & ({}^E \Omega(\mathcal{T}), D, [,]_{SN}) \\
 \downarrow \text{}^E L_{mod} & & \downarrow L_{mod} \\
 ({}^E A_*, 0) & \xrightarrow{\lambda_A} & ({}^E \Omega(\mathcal{A}), D),
 \end{array}
 \tag{4.1}$$

$$\begin{array}{ccc}
 ({}^E \Omega(\mathcal{D}), D + \partial, [,]_G) & \xleftarrow{\lambda_D} & ({}^E D_{poly}^*, \partial, [,]_G) \\
 \downarrow \mathcal{R}_{mod} & & \downarrow \text{}^E R_{mod} \\
 ({}^E \Omega(\mathcal{J}), D + \mathfrak{b}) & \xleftarrow{\lambda_C} & ({}^E C_*^{poly}, \mathfrak{b}),
 \end{array}$$

where the horizontal arrows correspond to embeddings of the sheaves of DGLAs (resp. of DGLA modules) constructed in the previous section. These embeddings are quasi-isomorphisms by theorems 3.4, 3.8 and propositions 3.5, 3.7.

Next, due to claims 1 and 2 in theorem 2.5 we have a fiberwise L_∞ -quasi-isomorphism

$$\mathcal{K} : ({}^E \Omega(\mathcal{T}), 0, [,]_{SN}) \xrightarrow{\sim} ({}^E \Omega(\mathcal{D}), \partial, [,]_G)
 \tag{4.2}$$

from the sheaf of DGLAs $({}^E\Omega(\mathcal{T}), 0, [,]_{SN})$ to the sheaf of DGLAs $({}^E\Omega(\mathcal{D}), \partial, [,]_G)$. Composing L_∞ -quasi-isomorphism (4.2) with the action of ${}^E\Omega(\mathcal{D})$ on ${}^E\Omega(\mathcal{J})$ we get an L_∞ -module structure on ${}^E\Omega(\mathcal{J})$ over ${}^E\Omega(\mathcal{T})$.

Due to claims 1 and 2 in theorem 2.6 we have a fiberwise L_∞ -quasi-isomorphism

$$(4.3) \quad \mathcal{S} : ({}^E\Omega(\mathcal{J}), \mathfrak{b}) \succ \rightarrow ({}^E\Omega(\mathcal{A}), 0)$$

from the sheaf of L_∞ -modules ${}^E\Omega(\mathcal{J})$ to the sheaf of DGLA modules ${}^E\Omega(\mathcal{A})$ over ${}^E\Omega(\mathcal{T})$.

Thus we get the following commutative diagram

$$(4.4) \quad \begin{array}{ccc} ({}^E\Omega(\mathcal{T}), 0, [,]_{SN}) & \xrightarrow{\mathcal{K}} & ({}^E\Omega(\mathcal{D}), \partial, [,]_G) \\ \downarrow \mathcal{L}_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^E\Omega(\mathcal{A}), 0) & \xleftarrow{\mathcal{S}} & ({}^E\Omega(\mathcal{J}), \mathfrak{b}), \end{array}$$

where by commutativity we mean that \mathcal{S} is a L_∞ -morphism of the sheaves of L_∞ -modules $({}^E\Omega(\mathcal{J}), \mathfrak{b})$ and $({}^E\Omega, 0)$ over the sheaf of DGLAs $({}^E\Omega(\mathcal{T}), 0, [,]_{SN})$ and the L_∞ -module structure on $({}^E\Omega(\mathcal{J}), \mathfrak{b})$ over $({}^E\Omega(\mathcal{T}), 0, [,]_{SN})$ is obtained by composing the L_∞ -morphism \mathcal{K} with the action \mathcal{R} (see 3.4 in [10]) of $({}^E\Omega(\mathcal{D}), \partial, [,]_G)$ on $({}^E\Omega(\mathcal{J}), \mathfrak{b})$.

Let us now restrict ourselves to an open subset $V \subset M$ such that $E|_V$ is trivial. Over any such subset the E -de Rham differential (1.5) is well defined for either of the sheaves ${}^E\Omega(\mathcal{A})$, ${}^E\Omega(\mathcal{T})$, ${}^E\Omega(\mathcal{J})$, and ${}^E\Omega(\mathcal{D})$. Furthermore, since the L_∞ -quasi-isomorphisms (4.2) and (4.3) are fiberwise we can add to all the differentials in diagram (4.4) the E -de Rham differential (1.5). Thus we get a new commutative diagram

$$(4.5) \quad \begin{array}{ccc} ({}^E\Omega(\mathcal{T})|_V, {}^E d, [,]_{SN}) & \xrightarrow{\mathcal{K}} & ({}^E\Omega(\mathcal{D})|_V, {}^E d + \partial, [,]_G) \\ \downarrow \mathcal{L}_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^E\Omega(\mathcal{A})|_V, {}^E d) & \xleftarrow{\mathcal{S}} & ({}^E\Omega(\mathcal{J})|_V, {}^E d + \mathfrak{b}) \end{array}$$

of the L_∞ -morphism \mathcal{K} and the morphism of L_∞ -modules \mathcal{S} .

We claim that

Proposition 4.1. *The L_∞ -morphism \mathcal{K} and the morphism of L_∞ -modules \mathcal{S} in (4.5) are quasi-isomorphisms.*

Proof. This statement follows easily from the standard argument of the spectral sequence. Indeed, we can naturally regard ${}^E\Omega(\mathcal{T})$ and ${}^E\Omega(\mathcal{D})$ (resp. ${}^E\Omega(\mathcal{J})$ and ${}^E\Omega(\mathcal{A})$) as sheaves of double complexes and the exterior degree provides us with the following descending filtration

$$F^p({}^E\Omega^*(\mathcal{B})) = \bigoplus_{k \geq p} {}^E\Omega^k(\mathcal{B}),$$

where \mathcal{B} is either \mathcal{T} or \mathcal{D} (resp. \mathcal{J} or \mathcal{A}).

The corresponding versions of Vey's [36] and Hochschild-Kostant-Rosenberg-Connes-Teleman [4], [19], [34] theorems for \mathbb{R}_{formal}^d imply that \mathcal{K} (resp. \mathcal{S}) induces a quasi-isomorphism on the level of E_0 . Hence, \mathcal{K} (resp. \mathcal{S}) induces a

quasi-isomorphism on the level of E_∞ . The standard snake lemma argument of homological algebra implies that \mathcal{K} (resp. \mathcal{S}) in (4.5) is a quasi-isomorphism. \square

On the open subset V we can represent the Fedosov differential (3.21) in the following (non-covariant) form

$$(4.6) \quad D = {}^E d + B \cdot ,$$

$$B = \sum_{p=0}^{\infty} \xi^i B_{i;j_1 \dots j_p}^k(x) y^{j_1} \dots y^{j_p} \frac{\partial}{\partial y^k} .$$

If we regard B as a section in ${}^E \Omega^1(V, \mathcal{T}^0)$ then the nilpotency condition $D^2 = 0$ says that B is a Maurer-Cartan section of the sheaf of DGLAs $({}^E \Omega(\mathcal{T}) \Big|_V, {}^E d, [,]_{SN})$. In the terminology of section 2 in [10] this means that the sheaf of DGLAs $({}^E \Omega(\mathcal{T}) \Big|_V, D, [,]_{SN})$ is obtained from $({}^E \Omega(\mathcal{T}) \Big|_V, {}^E d, [,]_{SN})$ via the twisting procedure by the Maurer-Cartan element B .

According to proposition 1 in section 2 of [10] the element

$$B_D = \sum_{k=1}^{\infty} \frac{1}{k!} \mathcal{K}_k(B, \dots, B)$$

is a Maurer-Cartan section of $({}^E \Omega(\mathcal{D}) \Big|_V, {}^E d + \partial, [,]_G)$. Moreover, due to claim 3 in theorem 2.5

$$B_D = B ,$$

where B is viewed as a section of the sheaf ${}^E \Omega^1(\mathcal{D}^0) \Big|_V$.

Thus twisting the L_∞ -quasi-isomorphism \mathcal{K} in (4.5) by the Maurer-Cartan element B we get the L_∞ -quasi-isomorphism

$$\mathcal{K}^{tw} : ({}^E \Omega(\mathcal{T}) \Big|_V, D, [,]_{SN}) \succrightarrow ({}^E \Omega(\mathcal{D}) \Big|_V, D + \partial, [,]_G) .$$

Since the DGLA module structure on ${}^E \Omega(\mathcal{A})$ over ${}^E \Omega(\mathcal{T})$ (resp. on ${}^E \Omega(\mathcal{J})$ over ${}^E \Omega(\mathcal{D})$) is honest the twist by the Maurer-Cartan element described in section 2 of [10] do not change these structures. Hence, by virtue of propositions 3 and 4 in [10] the twisting procedure turns diagram (4.5) into the commutative diagram

$$(4.7) \quad \begin{array}{ccc} ({}^E \Omega(\mathcal{T}) \Big|_V, D, [,]_{SN}) & \xrightarrow{\mathcal{K}^{tw}} & ({}^E \Omega(\mathcal{D}) \Big|_V, D + \partial, [,]_G) \\ \downarrow L_{mod} & & \downarrow R_{mod} \\ ({}^E \Omega(\mathcal{A}) \Big|_V, D) & \xleftarrow{\mathcal{S}^{tw}} & ({}^E \Omega(\mathcal{J}) \Big|_V, D + \mathfrak{b}) \end{array}$$

where \mathcal{S}^{tw} is a L_∞ -quasi-isomorphism obtained from \mathcal{S} by twisting via the Maurer-Cartan section B of the sheaf of DGLAs $({}^E \Omega(\mathcal{T}) \Big|_V, {}^E d, [,]_{SN})$.

We claim that the L_∞ -morphism \mathcal{K}^{tw} (resp. \mathcal{S}^{tw}) does not depend on the choice of the trivialization of E over V and hence is a well-defined L_∞ -morphism of sheaves of DGLAs (resp. sheaves of DGLA modules). Indeed, the term in B that depends on the choice of the trivialization of E is linear in the fiber coordinates y^i . But due to claim 4 in theorem 2.5 and claim 3 in theorem 2.6 this term contribute neither to \mathcal{K}^{tw} nor to \mathcal{S}^{tw} .

Thus the L_∞ -quasi-isomorphisms \mathcal{K}^{tw} and \mathcal{S}^{tw} are well defined and we arrive at the following commutative diagram

$$(4.8) \quad \begin{array}{ccc} ({}^E\Omega(\mathcal{T}), D, [\cdot, \cdot]_{SN}) & \xrightarrow{\mathcal{K}^{tw}} & ({}^E\Omega(\mathcal{D}), D + \partial, [\cdot, \cdot]_G) \\ \downarrow L_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^E\Omega(\mathcal{A}), D) & \xleftarrow{\mathcal{S}^{tw}} & ({}^E\Omega(\mathcal{J}), D + \mathfrak{b}). \end{array}$$

Assembling diagrams (4.1) and (4.8) we get the desired chain (2.3) of quasi-isomorphisms between the sheaves of DGLA modules $({}^E T_{poly}^*, {}^E A_*)$ and $({}^E D_{poly}^*, {}^E C_*^{poly})$. It is obvious from the construction that the terms and the quasi-isomorphisms of the resulting diagram (2.3) are functorial in the pair (E, ∂^E) , where ∂^E is a torsion-free connection on E . Thus, theorem 2.2 is proved. \square

4.2. Applications of the formality theorem. The obvious applications of the formality theorem for E -chains are related to the deformations associated with Poisson Lie algebroids. Namely, theorem 2.2 allows us to get an elegant description of the Hochschild homology and the traces of these deformations.

First, we recall that

Definition 4.2. *A Lie algebroid (E, M, ρ) equipped with an E -bivector $\pi \in \Gamma(M, {}^E T_{poly}^1)$ satisfying the Jacobi identity*

$$(4.9) \quad [\pi, \pi]_{SN} = 0$$

is called a Poisson Lie algebroid.

Following [26] a quantization of a Poisson Lie algebroid is a construction of an element

$$(4.10) \quad \Pi \in \Gamma(M, {}^E D_{poly}^1)[[\hbar]]$$

satisfying the condition of the classical limit

$$(4.11) \quad \Pi = 1 \otimes 1 \bmod \hbar, \quad \Pi - t(\Pi) = \hbar \pi \bmod \hbar,$$

and the ‘‘associativity’’ condition

$$(4.12) \quad [\Pi, \Pi]_G = 0.$$

Here \hbar is an auxiliary variable and t denotes the (cyclic) permutation of components of $\Pi \in \Gamma(M, {}^E D_{poly}^1)[[\hbar]] = \Gamma(M, \mathcal{U}E \otimes \mathcal{U}E)[[\hbar]]$.

Furthermore, two deformations Π and Π' of (E, M, ρ, π) are called *equivalent* if there exists a formal power series

$$\Psi = 1 + \hbar \Psi_1 + \hbar^2 \Psi_2 + \dots \in \Gamma(M, \mathcal{U}E)[[\hbar]]$$

such that

$$(4.13) \quad (\Delta \Psi) \Pi' = \Pi (\Psi \otimes \Psi),$$

where Δ is the coproduct (1.13) in $\mathcal{U}E$.

Thanks to the formality theorem for the sheaf of DGLAs ${}^E D_{poly}^*$ proved in [1] we have a bijective correspondence between the moduli spaces of Maurer-Cartan elements of the DGLA $\hbar \Gamma(M, {}^E T_{poly}^*)[[\hbar]]$ of E -polyvector fields and the DGLA

$\hbar\Gamma(M, {}^E D_{poly}^*)[[\hbar]]$ of E -polydifferential operators. In other words, if we consider the cone

$$(4.14) \quad \begin{aligned} \pi_{\hbar} &= \hbar\pi + \hbar^2\pi_1 + \hbar^3\pi_2 + \dots, \\ [\pi_{\hbar}, \pi_{\hbar}]_{SN} &= 0, \\ \pi_i &\in \Gamma(M, {}^E T_{poly}^1) \end{aligned}$$

of formal power series in \hbar acted upon by the Lie algebra $\hbar\Gamma(M, E)[[\hbar]]$

$$(4.15) \quad \pi_{\hbar} \rightarrow [u, \pi_{\hbar}], \quad u \in \hbar\Gamma(M, E)[[\hbar]],$$

then

Corollary 4.3. *The deformations (4.10) associated with a Poisson Lie algebroid (E, M, ρ, π) modulo the relation (4.13) are in a bijective correspondence with the points of the cone (4.14) modulo the action (4.15) of the pronilpotent group corresponding to the Lie algebra $\hbar\Gamma(M, E)[[\hbar]]$. \square*

An orbit $[\pi_{\hbar}]$ on the cone (4.14) corresponding to a deformation Π (4.10) is called *the class of the deformation* and any point π_{\hbar} of this orbit is called *a representative* of the class.

Given a deformation Π (4.10) associated with a Poisson Lie algebroid (E, M, ρ, π) one can define Hochschild chain complex of this deformation as the graded vector space

$$(4.16) \quad \Gamma(M, {}^E C_*^{poly})[[\hbar]]$$

equipped with the differential

$${}^E R_{\Pi} : {}^E C_*^{poly} \rightarrow {}^E C_{*+1}^{poly}.$$

Furthermore, one defines the Hochschild cochain complex of the deformation Π as the graded vector space

$$(4.17) \quad \Gamma(M, {}^E D_{poly}^*)[[\hbar]]$$

equipped with the differential

$$[\Pi,] : {}^E D_{poly}^* \rightarrow {}^E D_{poly}^{*+1}.$$

Due to claim 5 of proposition 2 in [10], claim 5 of proposition 3 in [10], and theorem 2.2 we get the following result:

Corollary 4.4. *Let Π be a deformation associated with a Poisson Lie algebroid (E, M, ρ, π) and let π_{\hbar} be a representative of the class of this deformation. Then the complex of Hochschild cohomology (4.17) of the deformation Π is quasi-isomorphic to the complex of E -polyvector fields*

$$(4.18) \quad (\Gamma(M, {}^E T_{poly}^*)[[\hbar]], [\pi_{\hbar},])$$

with the differential $[\pi_{\hbar},]$. The complex of Hochschild homology (4.16) of the deformation Π is quasi-isomorphic to the complex of E -forms

$$(4.19) \quad ({}^E \Omega(M)[[\hbar]], {}^E L_{\pi_{\hbar}})$$

with the differential ${}^E L_{\pi_{\hbar}}$. \square

Given a deformation Π (4.10) associated with a Poisson Lie algebroid (E, M, ρ, π) one can define a *trace*⁷ of the deformation Π as an $\mathbb{R}[[\hbar]]$ -linear functional

$$(4.20) \quad tr : \mathcal{O}(M)[[\hbar]] \rightarrow \mathbb{R}[[\hbar]]$$

satisfying the following condition

$$(4.21) \quad tr(j(\Pi) - j(t(\Pi))) = 0, \quad \forall j \in \Gamma(M, {}^E J_1^{poly}) \cap \ker \nabla^G.$$

It is not hard to see that corollary 4.4 implies the following statement:

Corollary 4.5. *Let Π be a deformation associated with a Poisson Lie algebroid (E, M, ρ, π) and let π_{\hbar} be a representative of the class of this deformation. Then the vector space of traces of the deformation Π is isomorphic to the vector space of continuous $\mathbb{R}[[\hbar]]$ -linear $\mathbb{R}[[\hbar]]$ -valued functionals on $\mathcal{O}(M)[[\hbar]]$ vanishing on all functions $f \in \mathcal{O}(M)[[\hbar]]$ of the following form*

$$f = j(\pi_{\hbar}), \quad j \in \Gamma(M, {}^E J_1^{poly}) \cap \ker \nabla^G,$$

where π_{\hbar} is viewed as a series E -bidifferential operators. \square

5. FORMALITY THEOREMS FOR HOLOMORPHIC LIE ALGEBROIDS

Let now M be a complex manifold. Let us write $T_{\mathbb{C}}M = T^{1,0} \oplus T^{0,1}$ for the decomposition of the (complexified) tangent bundle as the sum of the holomorphic tangent bundle and anti-holomorphic tangent bundle. We denote by \mathcal{O}_M the structure sheaf of holomorphic functions on M and by z^α local coordinates on M . We have to adapt the definition of holomorphic Lie algebroids:

Definition 5.1. *A holomorphic Lie algebroid over a complex manifold M is a holomorphic vector bundle E of finite rank whose sheaf of sections is a sheaf of Lie algebras equipped with a \mathcal{O}_M -linear map of sheaves of (complex) Lie algebras*

$$\rho : E \rightarrow T^{1,0},$$

satisfying the same condition described (for the smooth case) in formula (1.1).

Remark. This notion is different from the one of a complex Lie algebroid that we introduced in the remark of subsection 2.2.

5.1. Algebraic structures and the main theorem. Let E be a holomorphic Lie algebroid. As in section 1, one can define the following sheaves (which are also holomorphic vector bundles):

- ${}^E T_{poly}^*$ is the sheaf of E -polyvector fields. We regard ${}^E T_{poly}^*$ as a sheaf of DGLAs with the vanishing differential and with the Lie bracket $[\cdot, \cdot]_{SN}$ defined as in (1.3), (1.4).
- ${}^E A_*$ is the sheaf of E -differential forms with converted grading:

$$(5.1) \quad {}^E A_* = \wedge^{-*} E^\vee, \quad {}^E A_0 = \mathcal{O}_M.$$

We regard ${}^E A_*$ as a sheaf of DGLA modules over ${}^E T_{poly}^*$ with the vanishing differential and with the action ${}^E L$ defined as in (1.6).

⁷This notion is very important for formulations of various versions of algebraic index theorems for deformations associated with Poisson Lie algebroids [26].

- ${}^E D_{poly}^*$ is a sheaf of E -polydifferential operators. We regard ${}^E D_{poly}^*$ as a sheaf of DGLAs with the bracket $[\cdot]_G$ and the differential ∂ defined as in (1.24) and (1.25). Notice that the tensor product of sections (over \mathcal{O}_M) of ${}^E D_{poly}^*$ turns ${}^E D_{poly}^*$ into a sheaf of DGAs.
- ${}^E J_*^{poly}$ is the sheaf of E -polyjets

$${}^E J^{poly} = \bigoplus_{k \geq 0} {}^E J_k^{poly}, \quad {}^E J_*^{poly} := Hom_{\mathcal{O}_M}({}^E D_{poly}^*, \mathcal{O}_M),$$

which we regard as a sheaf of DGLA modules over ${}^E D_{poly}^*$ with the action ${}^E S$ and the differential \mathfrak{b} defined as in (1.31) and (1.35). The sheaf ${}^E J_*^{poly}$ is also equipped with the Grothendieck connection

$$(5.2) \quad \nabla^G : T^{1,0} \otimes {}^E J_*^{poly} \mapsto {}^E J_*^{poly}, \quad \nabla_u^G(j)(P) := \rho(u)(j(P)) - j(u \bullet P),$$

where $u \in \Gamma(U, T^{1,0})$ is a holomorphic vector field, $P \in \Gamma(U, {}^E D_{poly}^k)$, $j \in \Gamma(U, {}^E J_k^{poly})$ and the operation \bullet is defined in (1.23). The connection (5.2) is compatible the DGLA module structure on ${}^E J_*^{poly}$.

- ${}^E C_*^{poly}$ is the graded sheaf of ∇^G -flat E -polyjets with converted grading

$$(5.3) \quad {}^E C_*^{poly} := \ker \nabla^G \cap {}^E J_{-*}^{poly}.$$

Due to the compatibility of the Grothendieck connection (5.2) with the DGLA module structure on E -polyjets ${}^E C_*^{poly}$ can be viewed as a sheaf of DG modules over sheaf of DGLAs ${}^E D_{poly}^*$. We refer to ${}^E C_*^{poly}$ as a sheaf of Hochschild E -chains or E -chains for short.

The main result of this section can be formulated as follows:

Theorem 5.2. *For any holomorphic Lie algebroid E over a complex manifold M the sheaves of DGLA modules $({}^E T_{poly}^*, {}^E A_*)$ and $({}^E D_{poly}^*, {}^E C_*^{poly})$ are quasi-isomorphic.*

Omitting the sheaves of DGLA modules ${}^E A_*$ and ${}^E C_*^{poly}$ in the above theorem we get the following corollary:

Corollary 5.3. *For any holomorphic Lie algebroid E over a complex manifold M the sheaves of DGLAs ${}^E T_{poly}^*$ and ${}^E D_{poly}^*$ are quasi-isomorphic.*

We would like to mention that this corollary is parallel to the result of A. Yekutieli [39], who proved this statement for the tangent Lie algebroid $TM \rightarrow M$ of any smooth algebraic variety over a field \mathbb{K} for which $\mathbb{R} \subset \mathbb{K}$.

Notice that applying theorem 5.2 to the tangent algebroid $T^{1,0}M \rightarrow M$ we prove the following version of Tsygan's formality conjecture for complex manifolds:

Theorem 5.4. *For any complex manifold M the sheaf of DGLA modules $C^{poly}(M)$ of Hochschild chains over the sheaf $D_{poly}(M)$ of (holomorphic) polydifferential operators is formal. \square*

The proof of theorem 5.2 occupies the rest of the section.

5.2. Fedosov resolutions. First, we observe that any holomorphic Lie algebroid E can be viewed as a complex Lie algebroid in the sense of the remark in subsection 2.2, where the anchor map is naturally extended to the smooth sections of E . It is clear that the sheaf of Lie algebras $T^{0,1}$ acts on E and that this action commutes with ρ as ρ is holomorphic. Thus we get

Proposition 5.5. *Let F be the smooth vector bundle $F = E \oplus T^{0,1}$. Then F is a complex Lie algebroid over M with the anchor map $\rho_F : F \rightarrow T^{1,0} \oplus T^{0,1}$ given by $\rho_F \Big|_E = \rho$ and $\rho_F \Big|_{T^{0,1}} = \text{id} : T^{0,1} \rightarrow T^{0,1}$. \square*

For a holomorphic vector bundle \mathcal{B} over M we consider the sheaf of smooth F -differential forms with values in \mathcal{B} :

$$(5.4) \quad \begin{aligned} F\Omega(\mathcal{B}) &= \bigoplus_{p,q} F\Omega^{p,q}(\mathcal{B}), \\ F\Omega^{p,q}(\mathcal{B}) &= \wedge^p E^\vee \otimes \wedge^q T^{*0,1}M \otimes \mathcal{B} \end{aligned}$$

For sections

$$(5.5) \quad \begin{aligned} a &= \sum_{p,q} a_{i_1 \dots i_p; \alpha_1, \dots, \alpha_q}(z, \bar{z}) \xi^{i_1} \dots \xi^{i_p} d\bar{z}^{\alpha_1} \dots d\bar{z}^{\alpha_q}, \\ a_{i_1 \dots i_p; \alpha_1, \dots, \alpha_q}(z, \bar{z}) &\in \Gamma^{\text{smooth}}(U, \mathcal{B}) \end{aligned}$$

of $F\Omega(\mathcal{B})$ we reserve the local basis $\{\xi^i\}$ of anti-commuting fiber coordinates on E and the local basis $\{d\bar{z}^\alpha\}$ of antiholomorphic exterior forms on M . We denote by \bar{d} the Dolbeault differential

$$(5.6) \quad \bar{d} = d\bar{z}^\alpha \partial_{\bar{z}^\alpha} : F\Omega^{p,*}(\mathcal{B}) \mapsto F\Omega^{p,*+1}(\mathcal{B}).$$

It is obvious that the (DG) algebraic structures on the sheaves ${}^E T_{poly}^*$, ${}^E A_*$, ${}^E D_{poly}^*$, and ${}^E J_*^{poly}$, can be naturally extended to the sheaves ${}^F \Omega^{0,*}({}^E T_{poly}^*)$, ${}^F \Omega^{0,*}({}^E A_*)$, ${}^F \Omega^{0,*}({}^E D_{poly}^*)$, and ${}^F \Omega^{0,*}({}^E J_*^{poly})$. Similarly, the Grothendieck connection (5.2) on ${}^E J_*^{poly}$ extends to the operator

$$(5.7) \quad \nabla^G : T^{1,0} \otimes {}^F \Omega^{0,*}({}^E J_*^{poly}) \mapsto {}^F \Omega^{0,*}({}^E J_*^{poly}),$$

which is compatible with the action ${}^E S$ of ${}^F \Omega^{0,*}({}^E D_{poly}^*)$ on ${}^F \Omega^{0,*}({}^E J_*^{poly})$ and with the differential \mathfrak{b} on ${}^F \Omega^{0,*}({}^E J_*^{poly})$.

Since ${}^E T_{poly}^*$, ${}^E A_*$, ${}^E D_{poly}^*$, and ${}^E J_*^{poly}$ are holomorphic vector bundles it makes sense to speak about the Dolbeault differential (5.6)

$$(5.8) \quad \bar{d} : {}^F \Omega^{0,*}(\mathcal{B}) \mapsto {}^F \Omega^{0,*+1}(\mathcal{B}),$$

for \mathcal{B} being either ${}^E T_{poly}^*$, ${}^E A_*$, ${}^E D_{poly}^*$, or ${}^E J_*^{poly}$. It is obvious that \bar{d} is compatible with the (DG) algebraic structures on ${}^F \Omega^{0,*}(\mathcal{B})$ and with the Grothendieck connection (5.7) on ${}^F \Omega^{0,*}({}^E J_*^{poly})$.

Furthermore, due to the \bar{d} -Poincaré lemma we have

Proposition 5.6. *If \mathcal{B} is either ${}^E T_{poly}^*$, ${}^E A_*$, ${}^E D_{poly}^*$, or ${}^E J_*^{poly}$ then the canonical inclusion of sheaves*

$$(5.9) \quad \text{inc} : \mathcal{B} \hookrightarrow {}^F \Omega^{0,*}(\mathcal{B})$$

is a quasi-isomorphism of complexes of sheaves $(\mathcal{B}, 0)$ and $({}^F \Omega^{0,}(\mathcal{B}), \bar{d})$. The inclusion inc is compatible with the (DG) algebraic structures on \mathcal{B} , and ${}^F \Omega^{0,*}(\mathcal{B})$, and with the Grothendieck connection (5.2), (5.7). \square*

Due to this proposition it suffices to prove that the sheaves of DGLA modules $({}^F \Omega^{0,*}({}^E T_{poly}^*), {}^F \Omega^{0,*}({}^E A_*))$, and $({}^F \Omega^{0,*}({}^E D_{poly}^*), {}^F \Omega^{0,*}({}^E J_*^{poly}))$ are quasi-isomorphic. To show this we follow the lines of section 2 and introduce the formally completed symmetric algebra $\hat{S}(E^\vee)$ of the dual bundle E^\vee and (holomorphic) bundles \mathcal{T} , \mathcal{D} ,

\mathcal{A} , \mathcal{J} associated with $\hat{S}(E^\vee)$ (see page 15). As in section 2, \mathcal{T} and \mathcal{D} are sheaves of DGLAs while \mathcal{A} and \mathcal{J} are sheaves of DGLA modules over \mathcal{T} and \mathcal{D} , respectively. \mathcal{D} is also a sheaf of DGA's.

Next, we consider sheaves of smooth F -differential forms with values in the bundles $\hat{S}(E^\vee)$, \mathcal{T} , \mathcal{D} , \mathcal{A} , and \mathcal{J} . It is clear that the sheaves ${}^F\Omega(\hat{S}(E^\vee))$, ${}^F\Omega(\mathcal{A})$, ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{D})$, and ${}^F\Omega(\mathcal{J})$ acquire the corresponding (DG) algebraic structures and the Dolbeault differential (5.6) is obviously compatible with these structures.

Furthermore, we have the following obvious analogue of proposition 3.1

Proposition 5.7. *The sheaf ${}^F\Omega(\mathcal{T}^0)$ of F -forms with values in fiberwise vector fields is a sheaf of graded Lie algebras. The sheaves ${}^F\Omega(\hat{S}(E^\vee))$, ${}^F\Omega(\mathcal{A})$, ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{D})$, and ${}^F\Omega(\mathcal{J})$ are sheaves of modules over ${}^F\Omega(\mathcal{T}^0)$ and the action of ${}^F\Omega(\mathcal{T}^0)$ is compatible with the DG algebraic structures on ${}^F\Omega(\hat{S}(E^\vee))$, ${}^F\Omega(\mathcal{A})$, ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{D})$, ${}^F\Omega(\mathcal{J})$ and with the Dolbeault differential (5.6). \square*

Due to this proposition one can extend the following differential

$$\delta := \xi^i \frac{\partial}{\partial y^i} : {}^F\Omega^{*,q}(\hat{S}(E^\vee)) \rightarrow {}^F\Omega^{*+1,q}(\hat{S}(E^\vee))$$

of the sheaf of algebras ${}^F\Omega(\hat{S}(E^\vee))$ to the sheaves ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{D})$, ${}^F\Omega(\mathcal{A})$ and ${}^F\Omega(\mathcal{J})$ so that δ is compatible with the (DG) algebraic structures on ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{A})$, ${}^F\Omega(\mathcal{D})$, and ${}^F\Omega(\mathcal{J})$, and with the differential \bar{d} (5.6). Here $\{y^i\}$ (resp. $\{\xi^i\}$) denote commuting (resp. anticommuting) fiber coordinates of the bundle E .

We now have an analogue of proposition 3.2

Proposition 5.8. *For \mathcal{B} being either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} or \mathcal{D} and $q \geq 0$,*

$$H^{\geq 1}({}^F\Omega^{*,q}(\mathcal{B}), \delta) = 0.$$

Furthermore,

$$\begin{aligned} H^0({}^F\Omega^{*,q}(\hat{S}(E^\vee)), \delta) &\cong {}^F\Omega^{0,q}(M, \mathcal{O}_M), \\ (5.10) \quad H^0({}^F\Omega^{*,q}(\mathcal{A}_*), \delta) &\cong {}^F\Omega^{0,q}(M, {}^E A_*), \\ H^0({}^F\Omega^{*,q}(\mathcal{T}^*), \delta) &\cong {}^F\Omega^{0,q}(M, \wedge^{*+1}(E)) \end{aligned}$$

as sheaves of (graded) commutative algebras and

$$(5.11) \quad H^0({}^F\Omega^{*,q}(\mathcal{D}^*), \delta) \cong {}^F\Omega^{0,q}(M, \otimes^{*+1}(S(E)))$$

as sheaves of DGAs over \mathcal{O}_M .

Proof. As in proposition 3.2 it suffices to construct an operator ($q \geq 0$)

$$(5.12) \quad \kappa : {}^F\Omega^{*,q}(\mathcal{B}) \rightarrow {}^F\Omega^{*-1,q}(\mathcal{B})$$

such that for any section u of ${}^F\Omega(\mathcal{B})$ equation

$$(5.13) \quad u = \delta\kappa(u) + \kappa\delta(u) + \mathcal{H}(u),$$

is still true, where now

$$(5.14) \quad \mathcal{H}(u) = u \Big|_{y^i = \xi^i = 0}.$$

and y^i are as above fiber coordinates on E . As in the proof of proposition 3.2 we define κ on ${}^F\Omega(\hat{S}(E^\vee))$ by equation (3.12) and then extend it to ${}^F\Omega(\mathcal{T})$, ${}^F\Omega(\mathcal{A})$, and ${}^F\Omega(\mathcal{D})$ in the componentwise manner. \square

Let us choose a connection ∂^E on E which is compatible with the complex structure on E

$$(5.15) \quad \partial^E = {}^E d + \bar{d} + \xi^i \Gamma_i : {}^F \Omega^*(E) \rightarrow {}^F \Omega^{*+1}(E),$$

where $\xi^i \Gamma_i$ is locally a section of the sheaf ${}^E \Omega^1(\text{End}(E))$ and ${}^E d : {}^F \Omega_M^{*,q} \rightarrow {}^F \Omega_M^{*+1,q}$ is defined in (1.5).

It is not hard to show that such a connection always exists, and moreover, one can always choose ∂^E to be torsion free.

As in (3.16) we extend (5.15) to a derivation of the DG sheaves ${}^F \Omega(\hat{S}(E^\vee))$, ${}^F \Omega(\mathcal{A})$, ${}^F \Omega(\mathcal{T})$, ${}^F \Omega(\mathcal{D})$, and ${}^F \Omega(\mathcal{J})$:

$$(5.16) \quad \nabla = {}^E d + \Gamma \cdot + \bar{d} : {}^F \Omega^*(\mathcal{B}) \rightarrow {}^F \Omega^{*+1}(\mathcal{B}),$$

where \mathcal{B} is either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , \mathcal{D} , or \mathcal{J} , $\Gamma = -\xi^i \Gamma_{ij}^k y^j \frac{\partial}{\partial y^k}$, $\Gamma_{ij}^k(x)$ are Christoffel's symbols of the connection ∂^E (5.15) and $\Gamma \cdot$ denotes the action of Γ on the sections of the sheaves ${}^F \Omega(\mathcal{B})$. Due to proposition 5.7 ∇ (5.16) is compatible with the DG algebraic structures on ${}^F \Omega(\hat{S}(E^\vee))$, ${}^F \Omega(\mathcal{T})$, ${}^F \Omega(\mathcal{A})$, ${}^F \Omega(\mathcal{D})$, and ${}^F \Omega(\mathcal{J})$, and since ∇ is torsion free

$$(5.17) \quad \nabla \delta + \delta \nabla = 0.$$

Regarding (5.16) as a connection on \mathcal{B} one can see that the curvature of (5.16) has the components of type (2, 0) and (1, 1)

$$(5.18) \quad \nabla^2 = R^{2,0} + R^{1,1}, \quad R^{2,0} = ({}^E d + \Gamma)^2, \quad R^{1,1} = \bar{d} \Gamma.$$

We now prove the existence of a complex Fedosov differential D :

Theorem 5.9. *Let \mathcal{B} be either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , \mathcal{D} , or \mathcal{J} . There exists a section*

$$(5.19) \quad A = \sum_{s=2}^{\infty} \xi^k A_{k,i_1 \dots i_s}^j(z, \bar{z}) y^{i_1} \dots y^{i_s} \frac{\partial}{\partial y^j}$$

of the sheaf ${}^F \Omega^{1,0}(\mathcal{T}^0)$ and a section

$$(5.20) \quad \bar{A} = \sum_{s=2}^{\infty} d\bar{z}^\alpha \bar{A}_{\alpha,i_1 \dots i_s}^j(z, \bar{z}) y^{i_1} \dots y^{i_s} \frac{\partial}{\partial y^j}$$

of the sheaf ${}^F \Omega^{0,1}(\mathcal{T}^0)$ such that the derivation

$$(5.21) \quad D := \nabla - \delta + A \cdot + \bar{A} \cdot : {}^F \Omega^*(\mathcal{B}) \rightarrow {}^F \Omega^{*+1}(\mathcal{B})$$

is 2-nilpotent ($D^2 = 0$) and compatible with the DG algebraic structure on ${}^F \Omega(\mathcal{B})$.

Proof. Let us rewrite $D = D^{1,0} + D^{0,1}$ with

$$D^{1,0} = {}^E d + \Gamma \cdot - \delta + A \cdot, \quad D^{0,1} = \bar{d} + \bar{A} \cdot$$

and try to mimic the proof of theorem 3.3.

Due to (5.17) and (5.18) the condition $(D^{1,0})^2 = 0$ is equivalent to the equation

$$R^{2,0} + ({}^E d + \Gamma \cdot)A - \delta A + \frac{1}{2}[A, A]_{SN} = 0.$$

This equation has a solution obtained by iterations of the following equation (with respect to the degrees in fiber coordinates y_i 's)

$$A = \kappa R^{2,0} + \kappa(({}^E d + \Gamma \cdot)A + \frac{1}{2}[A, A]_{SN})$$

(the proof is the same as for theorem 3.3).

Using (5.18) once again we observe that the condition $D^{1,0}D^{0,1} + D^{0,1}D^{1,0} = 0$ is equivalent to

$$R^{1,1} + \bar{d}A + ({}^E d + \Gamma \cdot) \bar{A} - \delta \bar{A} + [A, \bar{A}]_{SN} = 0,$$

which, using the same arguments, has a solution obtained by iterations of the equation

$$\bar{A} = \kappa(R^{1,1} + \bar{d}A + ({}^E d + \Gamma \cdot) \bar{A} + [A, \bar{A}]_{SN}).$$

Indeed, denoting

$$C^{1,1} = R^{1,1} + \bar{d}A + ({}^E d + \Gamma \cdot) \bar{A} - \delta \bar{A} + [A, \bar{A}]_{SN},$$

and using that $\delta A = R^{2,0} + {}^E d + \Gamma \cdot A + \frac{1}{2}[A, A]_{SN}$ ($(D^{1,0})^2 = 0$), $\bar{d}R^{2,0} = 0$ and $\delta R^{1,1} = 0$ (Bianchi's identities for ∇) we get

$$({}^E d + \Gamma \cdot) C^{1,1} - \delta C^{1,1} + [A, C^{1,1}] = 0.$$

We have $\kappa C^{1,1} = 0$ by construction of \bar{A} and so, by the ‘‘Hodge-de Rham’’ decomposition (5.13), we have

$$C^{1,1} = \kappa(({}^E d + \Gamma \cdot) C^{1,1} + [A, C^{1,1}]).$$

The latter equation has the unique vanishing solution, which gives the result.

Let us now check the condition $(D^{0,1})^2 = 0$. This will be true if the section

$$C^{0,2} = \bar{d}\bar{A} + \frac{1}{2}[\bar{A}, \bar{A}] \in {}^F\Omega^{0,2}(\mathcal{T}^0)$$

is zero. One has again $D^{1,0}C^{0,2} = 0$ and $\kappa C^{0,2} = 0$ because it does not have ξ 's. As before, one can conclude that $C^{0,2} = 0$.

The compatibility of (5.21) with the corresponding DG algebraic structures follows from proposition 5.7. \square

We now describe the cohomology of the Fedosov differential D for the sheaves ${}^F\Omega(\hat{S}(E^\vee))$, ${}^F\Omega(\mathcal{A})$, ${}^F\Omega(\mathcal{T})$, and ${}^F\Omega(\mathcal{D})$

Theorem 5.10. *Let \mathcal{B} be either of the sheaves $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , or \mathcal{D} and $q \geq 0$. We have*

$$H({}^F\Omega^*(\mathcal{B}), D) \cong H({}^F\Omega^{0,*}(M, \mathcal{B}) \cap \ker \delta, \bar{d}).$$

as sheaves of (differential) graded (commutative) algebras.

Proof. Let us consider the double complex $({}^F\Omega^{*,*}(\mathcal{B}), D^{1,0} + D^{0,1})$. Using the degree in the fiber coordinates y^i we introduce on this complex a decreasing filtration. Applying the spectral sequence argument (as in the proof of theorem 3.4) and using proposition 5.8 we conclude that for any $i \geq 0$, the cohomology of the complex $({}^F\Omega^{*,i}(\mathcal{B}), D^{1,0})$ is concentrated in degree $*$ = 0. Therefore,

$$(5.22) \quad H({}^F\Omega^*(\mathcal{B}), D) = H({}^F\Omega^{0,*}(\mathcal{B}) \cap \ker D^{1,0}, D^{0,1}).$$

Following the lines of the proof of theorem 3.4 it is not hard to show that iterating the equation

$$(5.23) \quad \lambda(u) = u + \kappa(\nabla \lambda(u) + A \cdot \lambda(u) + \bar{A} \cdot \lambda(u)), \quad u \in {}^F\Omega^{0,q}(U, \mathcal{B}) \cap \ker \delta$$

we get an isomorphism of sheaves (of graded vector spaces)

$$(5.24) \quad \lambda : {}^F\Omega^{0,q}(\mathcal{B}) \cap \ker \delta \rightarrow {}^F\Omega^{0,q}(\mathcal{B}) \cap \ker D^{1,0},$$

and moreover, the map λ (5.24) has a natural inverse given by the map \mathcal{H} (5.14).

We claim that λ gives a quasi-isomorphism of complexes

$$\lambda : ({}^F\Omega^{0,*}(\mathcal{B}) \cap \ker \delta, \bar{d}) \rightarrow ({}^F\Omega^*(\mathcal{B}), D).$$

Indeed, due to (5.22) it suffices to show that for any $u \in {}^F\Omega^{0,q}(U, \mathcal{B}) \cap \ker \delta$, one has

$$\lambda(\bar{d}(u)) = D^{0,1}\lambda(u).$$

The term $\lambda(\bar{d}(u))$ is the only element in ${}^F\Omega^{0,q}(\mathcal{B})$ such that $\mathcal{H}(\lambda(\bar{d}(u))) = \bar{d}(u)$ and $D^{1,0}\lambda(\bar{d}(u)) = 0$. It is clear that $\mathcal{H}(D^{0,1}\lambda(u)) = \bar{d}(u)$ and one has

$$D^{1,0}D^{0,1}\lambda(u) = -D^{0,1}D^{1,0}\lambda(u) = 0,$$

since map λ (5.23) lands in $\ker D^{1,0}$.

The map λ (5.24) is compatible with the corresponding multiplications in $\hat{S}(E^\vee)$, \mathcal{A} , \mathcal{T} , or \mathcal{D} since so is the map \mathcal{H} (5.14). The theorem is proved. \square

It is not hard to prove the following analogue of proposition 3.5 :

Proposition 5.11. *The composition*

$$(5.25) \quad \mathcal{H}' = \nu \circ \mathcal{H} : {}^F\Omega^{0,*}(\mathcal{T}) \cap \ker D^{1,0} \rightarrow {}^F\Omega^{0,*}({}^E T_{poly}^*)$$

of the map

$$(5.26) \quad \mathcal{H} : {}^F\Omega^{0,*}(\mathcal{T}) \cap \ker D^{1,0} \rightarrow {}^F\Omega^{0,*}(\mathcal{T}) \cap \ker \delta$$

with the identification of the sheaves $\mathcal{T}^* \cap \ker \delta$ and ${}^E T_{poly}^* \cong \wedge^{*+1} E$

$$(5.27) \quad \nu : \mathcal{T}^* \cap \ker \delta \xrightarrow{\sim} {}^E T_{poly}^*$$

is an isomorphism of the sheaves of DGLAs

$$(5.28) \quad ({}^F\Omega^{0,*}(\mathcal{T}) \cap \ker D^{1,0}, D^{0,1}, [,]_{SN}) \cong ({}^F\Omega^{0,*}({}^E T_{poly}^*), \bar{d}, [,]_{SN})$$

The map

$$(5.29) \quad \mathcal{H} : {}^F\Omega^{0,*}(\mathcal{A}_*) \cap \ker D^{1,0} \rightarrow {}^F\Omega^{0,*}({}^E A_*)$$

is an isomorphism of the sheaves of DGLA modules

$$({}^F\Omega^{0,*}(\mathcal{A}_*) \cap \ker D^{1,0}, D^{0,1}) \cong ({}^F\Omega^{0,*}({}^E A_*), \bar{d})$$

over the sheaf of DGLAs (5.28). \square

Thanks to equation (5.22) this proposition implies that the map \mathcal{H}' gives a quasi-isomorphism of the sheaves of DGLAs $({}^F\Omega^*(\mathcal{T}), D, [,]_{SN})$ and $({}^F\Omega^{0,*}({}^E T_{poly}^*), \bar{d}, [,]_{SN})$.

Playing with the PBW theorem for the Lie algebroids (as we did in the proof of proposition 3.6) and with the cup product in the sheaves \mathcal{D} and ${}^E D_{poly}^*$ (see equation (3.42)) one can prove the following analogue of proposition 3.7

Proposition 5.12. *There exists an isomorphism of the sheaves of DGLAs*

$$(5.30) \quad \mu' : ({}^F\Omega^{0,*}({}^E D_{poly}^*), \bar{d}, [,]_G) \xrightarrow{\sim} ({}^F\Omega^{0,*}(\mathcal{D}) \cap \ker D^{1,0}, D^{0,1}, [,]_G),$$

which is compatible with the DGA structures on the sheaves ${}^F\Omega^{0,*}({}^E D_{poly}^*)$ and ${}^F\Omega^{0,*}(\mathcal{D})$. \square

Thanks to equation (5.22) this proposition implies that the map μ' (5.30) gives a quasi-isomorphism of the sheaves of DGLAs $({}^F\Omega^*(\mathcal{D}), D, [,]_G)$ and $({}^F\Omega^{0,*}({}^E D_{poly}^*), \bar{d}, [,]_G)$.

Let us consider the map

$$(5.31) \quad \gamma : {}^F\Omega^{0,q}(\mathcal{J}_*) \rightarrow {}^F\Omega^{0,q}({}^E J_*^{poly}), \quad \gamma(j)(P) = (\mu'(P))(j) \Big|_{y^i=0},$$

where $j \in {}^F\Omega^{0,q}(U, \mathcal{J}_k)$ and P is a holomorphic section of ${}^E D_{poly}^k$.

For this map we have the following obvious analogue of theorem 3.8

Theorem 5.13. *For any $q \geq 0$*

$$(5.32) \quad H^q({}^F\Omega^*(\mathcal{J}), D) = H^q({}^F\Omega^{0,*}(\mathcal{J}) \cap \ker D^{1,0}, D^{0,1}).$$

and the map γ (5.31) provides us with an isomorphism of the sheaves of DGLA modules

$$(5.33) \quad \gamma : {}^F\Omega^{0,*}(\mathcal{J}_*) \xrightarrow{\sim} {}^F\Omega^{0,*}({}^E J_*^{poly})$$

over the sheaf of DGLAs

$$({}^F\Omega^{0,*}(\mathcal{D}) \cap \ker D^{1,0}, D^{0,1}, [,]_G) \cong ({}^F\Omega^{0,*}({}^E D_{poly}^*), \bar{d}, [,]_G).$$

The map γ sends the component $D^{1,0}$ to the Grothendieck connection (5.7) and the component $D^{0,1}$ to the Dolbeault differential \bar{d} (5.6). \square

5.3. End of the proof of theorem 5.2. We have constructed the following honest (L_∞) -quasi-isomorphisms of the sheaves of DGLA modules

- $\lambda_T : ({}^F\Omega^{0,*}(M, {}^E T_{poly}^*), \bar{d}, [,]_{SN}) \rightarrow ({}^F\Omega(\mathcal{T}), D, [,]_{SN}),$
- $\lambda_A : ({}^F\Omega^{0,*}(M, {}^E A_*), \bar{d}) \rightarrow ({}^F\Omega(\mathcal{A}_*), D),$
- $\lambda_D : ({}^F\Omega^{0,*}(M, {}^E D_{poly}^*), \bar{d}, [,]_G) \rightarrow ({}^F\Omega(\mathcal{D}), D, [,]_G),$ and
- $\lambda_C : ({}^F\Omega^{0,*}(M, {}^E C_*^{poly}), \bar{d}) \rightarrow ({}^F\Omega(\mathcal{J}), D).$

Namely, the map λ_T is the inverse of \mathcal{H}' (5.25) the map λ_A is the inverse of \mathcal{H} (5.29) $\lambda_D = \mu'$ (5.30), and λ_C is composition of the identification (5.3) and the inverse of γ (5.31).

Our results can be summarized in the following commutative diagrams

$$(5.34) \quad \begin{array}{ccc} ({}^F\Omega^{0,q}(M, {}^E T_{poly}^*), \bar{d}, [,]_{SN}) & \xrightarrow{\lambda_T} & ({}^F\Omega(\mathcal{T}), D, [,]_{SN}) \\ \downarrow \mathcal{L}_{mod} & & \downarrow \mathcal{L}_{mod} \\ ({}^F\Omega^{0,q}(M, {}^E A_*), \bar{d}) & \xrightarrow{\lambda_A} & ({}^F\Omega(\mathcal{A}), D), \\ \\ ({}^F\Omega(\mathcal{D}), D + \partial, [,]_G) & \xleftarrow{\lambda_D} & ({}^F\Omega^{0,q}(M, {}^E D_{poly}^*), \bar{d} + \partial, [,]_G) \\ \downarrow \mathcal{R}_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^F\Omega(\mathcal{J}), D + \mathfrak{b}) & \xleftarrow{\lambda_C} & ({}^F\Omega^{0,q}(M, {}^E C_*^{poly}), \bar{d} + \mathfrak{b}), \end{array}$$

where the action ${}^E R$ is obtained from the action ${}^E S$ of ${}^F \Omega^{0,*}(M, {}^E D_{poly}^*)$ on ${}^F \Omega^{0,q}(M, {}^E J_*^{poly})$ via the identification (5.3).

Due to claims 1 and 2 in theorem 2.5 and claims 1 and 2 in theorem 2.6 we get the following commutative diagram

$$(5.35) \quad \begin{array}{ccc} ({}^F \Omega(\mathcal{T}), 0, [,]_{SN}) & \xrightarrow{\mathcal{K}} & ({}^F \Omega(\mathcal{D}), \partial, [,]_G) \\ \downarrow L_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^F \Omega(\mathcal{A}), 0) & \xleftarrow{\mathcal{S}} & ({}^F \Omega(\mathcal{J}), \mathfrak{b}), \end{array}$$

where by commutativity we mean that \mathcal{S} is a morphism of the sheaves of L_∞ -modules $({}^F \Omega(\mathcal{J}), \mathfrak{b})$ and $({}^F \Omega, 0)$ over the sheaf of DGLAs $({}^F \Omega(\mathcal{T}), 0, [,]_{SN})$ and the L_∞ -module structure on $({}^F \Omega(\mathcal{J}), \mathfrak{b})$ over $({}^F \Omega(\mathcal{T}), 0, [,]_{SN})$ is obtained by composing the L_∞ -isomorphism \mathcal{K} with the action \mathcal{R} (see 3.4 in [10]) of $({}^F \Omega(\mathcal{D}), \partial, [,]_G)$ on $({}^F \Omega(\mathcal{J}), \mathfrak{b})$.

Let us now restrict ourselves to an open subset $V \subset M$ such that $E|_V$ is trivial. Over any such subset the E -de Rham differential (1.5) is well defined for either of the sheaves ${}^F \Omega(\mathcal{A}), {}^F \Omega(\mathcal{T}), {}^F \Omega(\mathcal{J})$, and ${}^F \Omega(\mathcal{D})$. So again, we get a new commutative diagram

$$(5.36) \quad \begin{array}{ccc} ({}^F \Omega(\mathcal{T})|_V, {}^E d + \bar{d}, [,]_{SN}) & \xrightarrow{\mathcal{K}} & ({}^F \Omega(\mathcal{D})|_V, {}^E d + \bar{d} + \partial, [,]_G) \\ \downarrow L_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^F \Omega(\mathcal{A})|_V, {}^E d + \bar{d}) & \xleftarrow{\mathcal{S}} & ({}^F \Omega(\mathcal{J})|_V, {}^E d + \bar{d} + \mathfrak{b}) \end{array}$$

in which the L_∞ -morphism \mathcal{K} and the morphism of L_∞ -modules \mathcal{S} are quasi-isomorphisms.

On the open subset V we can represent the Fedosov differential (3.21) in the following (non-covariant) form

$$(5.37) \quad D = {}^E d + \bar{d} + B \cdot + \bar{B} \cdot,$$

$$B = \sum_{p=0}^{\infty} \xi^i B_{i;j_1 \dots j_p}^k(z, \bar{z}) y^{j_1} \dots y^{j_p} \frac{\partial}{\partial y^k},$$

and

$$\bar{B} = \sum_{p=0}^{\infty} d\bar{z}^\alpha \bar{B}_{\alpha;j_1 \dots j_p}^k(z, \bar{z}) y^{j_1} \dots y^{j_p} \frac{\partial}{\partial y^k},$$

where the z^α are local coordinates on M . If we regard $B + \bar{B}$ as a section of ${}^F \Omega^1(\mathcal{T}^0)|_V$ then the nilpotency condition $D^2 = 0$ says that $B + \bar{B}$ is a Maurer-Cartan section of the sheaf of DGLAs $({}^F \Omega(\mathcal{T})|_V, {}^E d + \bar{d}, [,]_{SN})$.

Thus applying the twisting procedures developed in section 2 of [10] and using claim 3 of theorem 2.5 we get the following commutative diagram

$$(5.38) \quad \begin{array}{ccc} ({}^E\Omega(\mathcal{T})|_V, D, [\cdot, \cdot]_{SN}) & \xrightarrow{\mathcal{K}^{tw}} & ({}^E\Omega(\mathcal{D})|_V, D + \partial, [\cdot, \cdot]_G) \\ \downarrow L_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^E\Omega(\mathcal{A})|_V, D) & \xleftarrow{S^{tw}} & ({}^E\Omega(\mathcal{J})|_V, D + \mathfrak{b}), \end{array}$$

in which \mathcal{K}^{tw} is a L_∞ -quasi-isomorphism of the sheaves of DGLAs and S^{tw} is a L_∞ -quasi-isomorphism of the sheaves of DGLA modules.

Due to claim 4 in theorem 2.5 and claim 3 in theorem 2.6 the quasi-isomorphisms do not depend on the trivialization of E over V .

Thus we constructed the following commutative diagram of sheaves of DGLAs, DGLA modules and their L_∞ -quasi-isomorphisms:

$$(5.39) \quad \begin{array}{ccc} ({}^E\Omega(\mathcal{T}), D, [\cdot, \cdot]_{SN}) & \xrightarrow{\mathcal{K}^{tw}} & ({}^E\Omega(\mathcal{D}), D + \partial, [\cdot, \cdot]_G) \\ \downarrow L_{mod} & & \downarrow \mathcal{R}_{mod} \\ ({}^E\Omega(\mathcal{A}), D) & \xleftarrow{S^{tw}} & ({}^E\Omega(\mathcal{J}), D + \mathfrak{b}), \end{array}$$

Combining the diagrams in (5.34), (5.39) together with the proposition 5.6 we see that the sheaves of DGLA modules $({}^E T_{poly}^*, {}^E A_*)$ and $({}^E D_{poly}^*, {}^E C_*^{poly})$ are connected by chain of quasi-isomorphisms. Thus, theorem 5.2 is proved. \square

6. CONCLUDING REMARKS

It would be interesting to prove the corresponding version of the algebraic index theorem [26], [32], which should relate a cyclic chain in the complex associated with a deformation Π (4.10) to its principal part and characteristic classes of the Lie algebroid (E, M, ρ) . It would be also interesting to investigate how other characteristic classes [6], [15], [22] of Lie algebroids could enter this picture.

Corollary 5.3 does not in general give a chain of quasi-isomorphisms between the DGLAs $\Gamma(M, {}^E T_{poly}^*)$ and $\Gamma(M, {}^E D_{poly}^*)$ of global sections. However, since the sheaves of smooth forms ${}^F\Omega^{0,*}({}^E T_{poly}^*)$ and ${}^F\Omega^{0,*}({}^E D_{poly}^*)$ are soft one could speculate about the deformations associated with E as about the Maurer-Cartan elements of the DGLA ${}^F\Omega^{0,*}(M, {}^E D_{poly}^*)[[\hbar]]$. Using the correspondence between the Dolbeault and Čech pictures one could relate these speculations to Kontsevich's algebroid picture of deformation quantization of algebraic varieties [21].

Finally, we think that the technique of mixed resolutions proposed by A. Yekutieli [39] could help us to prove Tsygan's formality conjecture for Hochschild chains of the structure sheaf of a smooth algebraic varieties over an arbitrary field of characteristic 0.

REFERENCES

- [1] D. Calaque, Formality for Lie algebroids, *Comm. Math. Phys.* **257** (2005), no. 3, 563-578.
- [2] A. Canas da Silva and A. Weinstein, *Geometric Models for Noncommutative algebras*, Berkeley Mathematics Lecture Notes, AMS Book, 1999.
- [3] A.S. Cattaneo, G. Felder, and L. Tomassini, From local to global deformation quantization of Poisson manifolds, *Duke Math. J.* **115** (2002), no. 2, 329-352.

- [4] A. Connes, Noncommutative differential geometry, IHES Publ. Math. **62** (1985), 257-360.
- [5] A. Connes, M. Flato, and D. Sternheimer, Closed star-products and cyclic cohomology, Lett. Math. Phys. **24** (1992), no. 1, 1-12.
- [6] M. Crainic, Differentiable and algebroid cohomology, van Est isomorphisms, and characteristic classes, Comment. Math. Helv. **78** (2003), no. 4, 681-721.
- [7] J. Cuntz, G. Skandalis, and B. Tsygan, Cyclic homology in non-commutative geometry. *Encyclopaedia of Mathematical Sciences*, 121. Operator Algebras and Non-commutative Geometry, II. Springer-Verlag, Berlin, 2004.
- [8] Yu. Daletski and B. Tsygan, Operations on cyclic and Hochschild complexes, Methods Func. Anal. Topology, **5** (1999), no. 4, 62-86.
- [9] V. Dolgushev, Covariant and equivariant formality theorems, Adv. Math. **191** (2005), no. 1, 147-177.
- [10] V. Dolgushev, A formality theorem for Hochschild chains, in Adv. Math. **200** (2005), no. 1, 51-101.
- [11] V. Dolgushev, A Proof of Tsygan's formality conjecture for an arbitrary smooth manifold, PhD thesis M.I.T., math.QA/0504420.
- [12] B. Fedosov, A simple geometrical construction of deformation quantization, J. Diff. Geom. **40** (1994), 213-238.
- [13] B. Fedosov, Deformation quantization and index theory, Akademie Verlag, Berlin, 1996.
- [14] G. Felder and B. Shoikhet, Deformation quantization with traces, Lett. Math. Phys. **53** (2000), no. 1, 75-86.
- [15] R.L. Fernandes, Lie Algebroids, Holonomy and Characteristic Classes, Adv. in Math. **170** (2002), no. 1, 119-179.
- [16] M. Gerstenhaber, The cohomology structure of an associative ring, Ann. Math. **78** (1963) 267-288.
- [17] E. Getzler, Cartan homotopy formulas and the Gauss-Manin connection in cyclic homology, in *Quantum deformations of algebras and their representations*, Israel Math. Conf. Proc. **7** (1993) 65-78.
- [18] V. Hinich and V. Schechtman, Homotopy Lie algebras, I.M. Gelfand Seminar, Adv. Sov. Math. **16** (1993), no. 2, 1-28.
- [19] G. Hochschild, B. Kostant, and A. Rosenberg, Differential forms on regular affine algebras, Trans. Amer. Math. Soc. **102** (1962), 383-408.
- [20] M. Kontsevich, Deformation quantization of Poisson manifolds, Lett. Math. Phys. **66** (2003), no. 3, 157-216.
- [21] M. Kontsevich, Deformation quantization of algebraic varieties. EuroConférence Moshé Flato 2000, Part III (Dijon). Lett. Math. Phys. **56** (2001), no. 3, 271-294.
- [22] S.L. Lyakhovich, A.A. Sharapov, Characteristic classes of gauge systems, Nucl. Phys. B703 (2004) 419-453.
- [23] K. Mackenzie, Lie groupoids and Lie algebroids in differential geometry, Lecture Notes Series **124**, London Mathematical Society, Cambridge University Press, 1987.
- [24] R. Mazzeo and R.B. Melrose, Pseudodifferential operators on manifolds with fibred boundaries, Asian J. Math. **2** (1998), 833-866.
- [25] B. Monthubert, Pseudodifferential calculus on manifolds with corners and groupoids, Proc. Amer. Mat. Soc. **127** (1999), no. 10, 2871-2881.
- [26] R. Nest and B. Tsygan, Deformations of symplectic Lie algebroids, deformations of holomorphic symplectic structures and index theorems, Asian J. Math. **5** (2001), no. 4, 599-633.
- [27] V. Nistor, A. Weinstein, and P. Xu, Pseudodifferential operators on differential groupoids, Pacific J. Math. **189** (1999) 117-152.
- [28] G.S. Rinehart, Differential forms on general commutative algebras, Trans. Amer. Math. Soc. **108** (1963) 195-222.
- [29] B. Shoikhet, A proof of Tsygan formality conjecture for chains, Adv. Math. **179** (2003), no. 1, 7-37.
- [30] B. Shoikhet, On the cyclic formality conjecture, math.QA/9903183.
- [31] D. Tamarkin, Another proof of M. Kontsevich formality theorem, math.QA/9803025.
- [32] D. Tamarkin and B. Tsygan, Cyclic formality and index theorems, Lett. Math. Phys. **56** (2001) 85-97.
- [33] D. Tamarkin, B. Tsygan, Noncommutative differential calculus, homotopy BV algebras and formality conjectures, Methods Funct. Anal. Topology **6** (2000), no.2, 85-100.

- [34] N. Teleman, Microlocalisation de l'homologie de Hochschild, C.R. Acad. Sci. Paris, t. 326, Serie I (1998) 1261-1264.
- [35] B. Tsygan, Formality conjectures for chains, in *Differential topology, infinite dimensional Lie algebras and applications*, 261-274, Amer. Math. Soc. Transl. Ser. 2 **194**, Amer. Math. Soc., Providence, RI, 1999.
- [36] J. Vey, Déformation du crochet de Poisson sur une variété symplectique, Comment. Math. Helv. **50** (1975), 421-454.
- [37] P. Xu, Quantum groupoids, Comm. Math. Phys. **206** (2001) 539-581.
- [38] E. Winkelkemper, The graph of a foliation, Ann. Global Anal. Geom. (1983) 51-75.
- [39] A. Yekutieli, Deformation Quantization in Algebraic Geometry, Adv. Math. **198** (2005), no.1, 383-432.

IRMA, 7 RUE RENÉ DESCARTES, F-67084 STRASBOURG, FRANCE

E-mail address: calaque@math.u-strasbg.fr

E-mail address: halbout@math.u-strasbg.fr

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139, USA

E-mail address: vald@math.mit.edu