

From Topological Strings to Integrable Hierarchies & back.

- Lecture # 2:
- * Tautological classes on $\bar{M}_{g,n}$
 - * Admissible G -covers
 - * G -Hodge classes
 - * Application: computing $\int_{H_g} \lambda_g \lambda_{g-1}$
 - * C.R.C. à la Bryan-Graber: $[\mathbb{C}^2/\mathbb{Z}_2]$

§1. Tautological Classes on $\bar{M}_{g,n}$

These are Chow classes on $\bar{M}_{g,n}$ constructed in a 'natural' geometric way. We focus on two families of such classes.

Ⓐ Ψ classes / Gravitational Descendants

$$\mathbb{L}_i = \mathbb{L}_i^*$$


$$\downarrow$$

$$\bar{M}_{g,n} \xleftarrow[\sigma_i]{\pi} \bar{M}_{g,n+1}$$

The i -th cotangent line bundle on $\bar{M}_{g,n}$:

$$\mathbb{L}_i := \sigma_i^*(\omega_\pi)$$

Fiber over a point $[(C, p_1, \dots, p_n)]$ is $T_{p_i}^* C$



Def: $\Psi_i := c_1(\mathbb{L}_i)$

Importance:

① In $g=0$ they are boundary class \Rightarrow give rise to T.R.R.

② If $D = \left[\begin{array}{c} X \\ p_1 = p_2 \end{array} \right] \Rightarrow N_{D/\bar{M}_{g,n}} \cong \mathbb{L}_1 \otimes \mathbb{L}_2$

(B) Hodge Classes

(2)

$$\begin{array}{c} \mathbb{E}_g \\ \downarrow \\ \bar{M}_g \leftarrow \bar{M}_{g,1} \end{array}$$

The Hodge bundle \mathbb{E}_g is a rank g vector bundle:

$$\mathbb{E}_g := \pi_* \omega_\pi$$

Fiber over $[C] \in \mathcal{M}_g$:

- $H^0(C, K_C)$: global sections of the canonical bundle;
- $\Omega^1(C)$: global holomorphic 1-forms on C ;
- $H^1(C, \mathcal{O}_C)^*$: by Serre duality.

Nice properties of \mathbb{E}_g :

$$\textcircled{1} \mathbb{E}_g|_{[\chi_{g_1, g_2}]} \cong \mathbb{E}_{g_1} \oplus \mathbb{E}_{g_2}$$

$$\textcircled{2} \mathbb{E}_g|_{[\alpha^{g-1}]} \cong \bar{\mathbb{E}}_{g-1} \oplus \mathcal{O}$$

\textcircled{3} (Mumford's Relations): $\mathbb{E} \oplus \mathbb{E}^\vee$ is a flat bundle
(\Rightarrow all Chern classes ($\neq c_0$) vanish)

Def: $\lambda_i := c_i(\mathbb{E}_g)$ - i th Hodge class.

Remark: properties above translate to combinatorial relations:

$$\textcircled{1} \Rightarrow \lambda_g \lambda_{g-1} |_{\bar{M}_g - \mathcal{M}_g} \equiv 0$$

$$\textcircled{2} \Rightarrow \lambda_g |_{[\alpha]} \equiv 0$$


$$\textcircled{3} \Rightarrow \lambda_g^2 = 0 \quad g \neq 0$$

$$\lambda_2 = \frac{\lambda_1^2}{2}$$

§2. Admissible G-covers

(3)

I want to describe a moduli space for:



- nodes correspond to nodes
- Kissing condition: matching ramification profile at the shadows of the nodes.

- Fixing ramification profiles is equivalent to specifying image of the (refined) evaluation maps to IBG.

Such a creature is called an admissible G-cover

Abramovich-Corti-Vistoli:

{ Stack of Admissible
G-covers + assigned
monodromy types }

← EQUIV. →

{ Component of $\bar{\mathcal{M}}_{0,n}(\text{BG}, 0)$
cut out by the class
 $\prod_i \text{ev}_i^*([1_{g_i}])$ }

§3. Tautological classes on Admissible Covers

(A) Ψ classes

Ψ classes are defined similarly. Only caveat, there are 2 types:

- stacky Ψ classes
- coarse Ψ classes

[1onel]: $\boxed{\text{stacky } \Psi_i = \frac{1}{|G_P|} \text{ coarse } \Psi_i}$

(B) Hodge classes:

$$\begin{array}{ccc}
 f^*IE_g & \longrightarrow & IE_g \\
 \downarrow & & \downarrow \\
 \bar{M}_{0,n}^{(BG)} & \xrightarrow{f} & \bar{M}_g \\
 \left[\begin{array}{c} E^{2g} \\ \downarrow \\ \mathbb{C} \end{array} \right] \rightarrow BG & \mapsto & [E]
 \end{array}$$

A 'rank' g Hodge bundle is obtained by pull back from \bar{M}_g .

$f^*IE_g := IE$ inherits a G -action from the action on the curves.

$$IE = \bigoplus_{\rho \text{ irreps of } G} IE^\rho \leftarrow \text{possibly many copies of } \rho$$

Def: $\lambda_i^\rho := c_i(IE^\rho)$ - G -Hodge classes.

Top intersection numbers of G -Hodge classes are called G -Hodge integrals.

Remark: G -Hodge classes enjoy combinatorial properties (vanishing, splitting along the boundary) similar to ordinary Hodge classes.

§4. Application: computing $\tan(\frac{x}{2})$ without GRR

Thm (Faber-Pandharipande)

$$H(x) := \sum_{g=1}^{\infty} \left(\int_{H_g} \lambda_g \lambda_{g-1} \right) \frac{x^{2g-1}}{(2g-1)!} = \frac{1}{2} \tan\left(\frac{x}{2}\right)$$

Proved in '90s using GRR.

COMBINATORIAL PROOF (Bertram, C-, Todorov):

① Setting up the 'right' generating functions

$$F(x) := \sum_{g=1}^{\infty} \left(\int_{H_g} \lambda_g \lambda_{g-1} \right) \frac{x^{2g+1}}{(2g+1)!} \quad F_i(x) := \sum_{g=i}^{\infty} \left(\int_{H_g} \lambda_g \lambda_{g-i} \psi^{i-1} \right) \frac{x^{2g+1}}{(2g+1)!} \quad \left(F_0(x) = \frac{x}{2} \right)$$

$$G(x) := F'(x)$$

$$G_i(x) := F_i'(x)$$

$$G_0 = \frac{1}{2}$$

$$H(x) := F''(x)$$


$$H_i(x) := F_i''(x)$$

② Auxiliary Localization Computations


(A) $\int_{M_{0, 2g+2}(\mathbb{P}^1 \times \mathbb{B}\mathbb{Z}_2, 1)}$ $\lambda_g \lambda_{g-2} \omega_1^*(0) \omega_2^*(0) \omega_3^*(\infty) = 0$ by dimension reasons.
twisted points

Compute 0 by localization:

Fixed loci:



$$\int_{H_g} \frac{\lambda_g \lambda_{g-2}}{1-\psi} = \int_{\bar{H}_g} \lambda_g \lambda_{g-2} \psi$$



STACKY GLUING

$$2 \binom{2g-1}{2g_2} \int_{H_{g_1}} \lambda_{g_1} \lambda_{g_1-1} \int_{H_{g_2}} \lambda_{g_2} \lambda_{g_2-1}$$

In terms of generating functions:

(6)

$$G'_2 = 2GG' \Rightarrow G_2 = G^2 = \frac{1}{2} \cdot \frac{2^2}{2!} G^2$$

(B) Repeat the trick:

$$\int \lambda_g \lambda_{g-3} = 0 \Rightarrow G_3 = \frac{1}{2} \frac{2^3}{3!} G^3$$

And in general:

$$\int \lambda_g \lambda_{g-i} = 0 \Rightarrow G_i = \frac{1}{2} \frac{2^i}{i!} G^i$$

And:

$$\boxed{\sum_0^\infty G_i = \frac{1}{2} e^{2G}} \quad (1)$$

(C) $\int_{\bar{M}_{0,2g+2}(\mathbb{P}^1 \times \mathbb{B}\mathbb{Z}_2, 1)} e^{e^g} (R^1 \pi_* f^*(\mathcal{O} \oplus \mathcal{O}(-1))) = 0$


Linearize bundles with weights

$$\begin{matrix} 0 & \infty \\ 0(-1) & \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} \\ 0 & \end{matrix}$$

Fixed loci:



$$\rightarrow (-1)^{g+1} \int_{\bar{H}_{g,1}} \psi^{2g} = \frac{1}{2}$$



$$\rightarrow (-1)^g \frac{2}{2} \binom{2g+2}{2g+1} \int_{\bar{H}_{g_1}} \psi^{2g, -1} \int_{\bar{H}_{g_2}} \lambda_g \lambda_{g-1} + \dots + \lambda_g \psi^{g-1}$$

In g.f. form: $\frac{1}{2} \sin x \sum_0^\infty F_i = \frac{1}{2} (1 - \cos x)$

$$\Rightarrow \boxed{\sum_0^\infty F_i = \tan \frac{x}{2}} \quad (2)$$

(3) Calculus exercise:

$$(1) + (2) \Rightarrow F = \frac{1}{2} \tan \left(\frac{x}{2} \right) \quad \square$$


§5. Crepant Resolution Conjecture à la Bryan-Graber

\mathcal{X} a Gorenstein orbifold ^{has a $K_{\mathcal{X}}$ line bundle} satisfying Hard Lefschetz and admitting a crepant resolution Y ^{$I: IB\mathcal{X} \rightarrow IB\mathcal{X}$ preserves age}
 $\pi^* K_{\mathcal{X}} = K_Y$

Then there is a graded linear ISO
 $L: H^*(Y) \rightarrow H^*_{CR}(\mathcal{X})$

such that:

- (1) L behaves naturally w.r.t. non exceptional classes
- (2) There is an analytic continuation of f^Y from $q_{exc} = 0$ to $q_{exc} = \vec{c}$ (\vec{c} a vector of roots of 1) such that:
- (3) $F^{\mathcal{X}} = F^Y$ after substituting $y = L(x)$ and specializing $q_{exc} = \vec{c}$



Example: the pair $[\mathbb{C}^2/\mathbb{Z}_2], \mathcal{O}_{\mathbb{P}^1}(-2)$

Ⓐ The GW theory of \mathcal{X}

- \mathcal{X} has no curve classes \Rightarrow only $\beta = 0$
- $I\mathcal{X} = \mathbb{C}^2/\mathbb{Z}_2 \amalg B\mathbb{Z}_2$
 age 0 age 1
 ψ ψ
 \perp -1

$$\langle (-1)^n \rangle_0 = \int_{[\overline{M}_{0,n}(\mathbb{C}^2/\mathbb{Z}_2, 0)]^{nr}} \prod e_i^*(-1) = \int_{[\overline{M}_{0,n}(B\mathbb{Z}_2, 0)]} \prod e_i^*(-1) \cdot e(O_b) \stackrel{n=2g+2}{=} \int_{H_g} \lambda_g \lambda_{g-1}$$

$$F^{\mathbb{R}} = \left\{ \begin{array}{l} \text{3-pt} \\ \text{inv's} \end{array} \right\} - \iiint \tan\left(\frac{x}{2}\right)$$

(B) GW theory of $\mathbb{P}^1 = \mathbb{C}P^1(-2)$

- Only 1 curve class (0-section)
- $H^*(Y) = H^0 \oplus H^2$
 $\quad \quad \quad \psi \quad \quad \psi$
 $\quad \quad \quad 1 \quad \quad pt$

DIVISOR EQ'N

$$F^Y = \left\{ \begin{array}{l} \text{3-pt} \\ \text{inv's} \\ \text{of deg 0} \end{array} \right\} + \sum_{d>0} \langle pt^n \rangle_d \frac{y^d}{d!} = \left\{ \quad \right\} + \sum_{d>0} \langle \phi \rangle_d e^{dy}$$

$\langle \phi \rangle_d$ - can be computed via localization (Aspinwall-Morrison formula)

$$\langle \phi \rangle_d = \frac{1}{d^3}$$

$$F^Y = \left\{ \begin{array}{l} \text{3-pt} \\ d=0 \end{array} \right\} + \sum_{d>0} \frac{1}{d^3} e^{dy}$$

(C) Change of variables

Set $y = ix$
 $q = -1$

Calculus exercise:

$$\left(\frac{d}{dx}\right)^3 F^Y(ix, q=-1) = \left(\frac{d}{dx}\right)^3 F^{\mathbb{R}}(x)$$